

B41

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization  
International Bureau(43) International Publication Date  
11 October 2001 (11.10.2001)

PCT

(10) International Publication Number  
**WO 01/74761 A1**

- (51) International Patent Classification<sup>7</sup>: **C07C 271/30**, 275/40, 271/58, 335/20, 235/84, C07D 277/42, A61K 31/65
- (74) Agents: **HANLEY, Elizabeth, A.** et al.; Lahive & Cockfield, LLP, 28 State Street, Boston, MA 02109 (US).
- (21) International Application Number: PCT/US01/10342
- (81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZW.
- (22) International Filing Date: 30 March 2001 (30.03.2001)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data:  
60/193,972 31 March 2000 (31.03.2000) US  
60/193,879 31 March 2000 (31.03.2000) US  
60/xxx,xxx 29 March 2001 (29.03.2001) US
- (84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).
- (71) Applicant: **TRUSTEES OF TUFTS COLLEGE** [US/US]; 4th floor, Ballou Hall, Medford, MA 02155 (US).
- (72) Inventors: **NELSON, Mark, L.**; 735 Worcester Road, Wellesley, MA (US). **LEVY, Stuart, B.**; 144 Warren Avenue, Boston, MA 02116 (US). **FRECHETTE, Roger**; 40 Estate Lane, Reading, MA 01867 (US). **BOWSER, Todd, E.**; 88 Haggerty Road, Charlton, MA 01507 (US). **ISMAIL, Mohamed, Y.**; 5 Hunt Road, Bedford, MA 01730 (US).
- Published:**  
— with international search report  
— before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments
- For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.*



WO 01/74761 A1

(54) Title: 7-AND 9-CARBAMATE, UREA, THIOUREA, THIOCARBAMATE, AND HETEROARYL-AMINO SUBSTITUTED TETRACYCLINE COMPOUNDS

(57) Abstract: Substituted tetracycline compounds, methods of synthesis, and methods of use are discussed. Tetracyclines useful for treating tetracycline related disorders are also discussed. Intermediates useful for synthesizing other tetracycline compounds are also included.

- 1 -

## **7- AND 9- CARBAMATE, UREA, THIOUREA, THIOCARBAMATE, AND HETEROARYL-AMINO SUBSTITUTED TETRACYCLINE COMPOUNDS**

### **Related Applications**

5           This application claims priority to U.S. Provisional Application No. 60/XXX,XXX, filed March 29, 2001, entitled "7- and 9- Carbamate, Urea, Thiourea, Thiocarbamate, and Heteroaryl-Amino Substituted Tetracycline Compounds;" U.S. Provisional Application No. 60/193,972, filed March 31, 2000, entitled "Methods for Synthesizing 7- or 9- Substituted Tetracycline Compounds and Reactive Intermediates;"  
10          and to U.S. Provisional Application No. 60/193,879, filed March 31, 2000, entitled "9- Substituted Tetracycline Compounds." The entire contents of all of the aforementioned applications are hereby incorporated herein by reference.

### **Background of the Invention**

15           The development of the tetracycline antibiotics was the direct result of a systematic screening of soil specimens collected from many parts of the world for evidence of microorganisms capable of producing bacteriocidal and/or bacteriostatic compositions. The first of these novel compounds was introduced in 1948 under the name chlortetracycline. Two years later, oxytetracycline became available. The  
20          elucidation of the chemical structure of these compounds confirmed their similarity and furnished the analytical basis for the production of a third member of this group in 1952, tetracycline. A new family of tetracycline compounds, without the ring-attached methyl group present in earlier tetracyclines, was prepared in 1957 and became publicly available in 1967.


25           Recently, research efforts have focused on developing new tetracycline antibiotic compositions effective under varying therapeutic conditions and routes of administration. New tetracycline analogues have also been investigated which may prove to be equal to or more effective than the originally introduced tetracycline compounds. Examples include U.S. Patent Nos. 3,957,980; 3,674,859; 2,980,584;  
30          2,990,331; 3,062,717; 3,557,280; 4,018,889; 4,024,272; 4,126,680; 3,454,697; and 3,165,531. These patents are representative of the range of pharmaceutically active tetracycline and tetracycline analogue compositions.

- 2 -

Historically, soon after their initial development and introduction, the tetracyclines were found to be highly effective pharmacologically against rickettsiae; a number of gram-positive and gram-negative bacteria; and the agents responsible for lymphogranuloma venereum, inclusion conjunctivitis, and psittacosis. Hence, tetracyclines became known as "broad spectrum" antibiotics. With the subsequent establishment of their *in vitro* antimicrobial activity, effectiveness in experimental infections, and pharmacological properties, the tetracyclines as a class rapidly became widely used for therapeutic purposes. However, this widespread use of tetracyclines for both major and minor illnesses and diseases led directly to the emergence of resistance to these antibiotics even among highly susceptible bacterial species both commensal and pathogenic (*e.g.*, *pneumococci* and *Salmonella*). The rise of tetracycline-resistant organisms has resulted in a general decline in use of tetracyclines and tetracycline analogue compositions as antibiotics of choice.

#### Summary of the Invention

The invention pertains, at least in part, to substituted tetracycline compounds of the formula (I):



(I)

wherein:

X is CHC(R<sup>13</sup>Y'Y), CR<sup>6</sup>R<sup>6</sup>, S, NR<sup>6</sup>, or O;

R<sup>2</sup> is hydrogen, alkyl, alkynyl, alkoxy, alkylthio, alkylsulfinyl, alkylsulfonyl, alkylamino, arylalkyl, aryl, heterocyclic, heteroaromatic or a prodrug moiety;

R<sup>4</sup> and R<sup>4'</sup> are each hydrogen, alkyl, alkenyl, alkynyl, alkoxy, alkylthio, alkylsulfinyl, alkylsulfonyl, alkylamino, arylalkyl, aryl, heterocyclic, heteroaromatic or a prodrug moiety;

R<sup>2'</sup>, R<sup>3</sup>, R<sup>10</sup>, R<sup>11</sup> and R<sup>12</sup> are each hydrogen or a pro-drug moiety;

R<sup>5</sup> is hydrogen, hydroxyl, or a prodrug moiety;

- 3 -

$R^6$ ,  $R^{6'}$ , and  $R^8$  are each independently hydrogen, alkyl, alkenyl, alkynyl, aryl, alkoxy, alkylthio, alkylsulfinyl, alkylsulfonyl, alkylamino, arylalkyl, or halogen;

$R^7$  is hydrogen, dialkylamino, heteroaryl-amino, or  $NR^{7c}C(=W')WR^{7a}$ ;

$R^{13}$  is hydrogen, hydroxy, alkyl; alkenyl; alkynyl; alkoxy; alkylthio; alkylsulfinyl; alkylsulfonyl; alkylamino; or an arylalkyl;

$Y'$  and  $Y$  are each independently hydrogen, halogen, hydroxyl, cyano, sulfhydryl, amino, alkyl, alkenyl, alkynyl, alkoxy, alkylthio, alkylsulfinyl, alkylsulfonyl, alkylamino, or arylalkyl;

$R^9$  is hydrogen, heteroaryl-amino, or  $NR^{9c}C(=Z')ZR^{9a}$ ;

$Z$  is  $CR^{9d}R^{9e}$ ,  $NR^{9b}$ , or O;

$Z'$  is O or S;

$R^{9a}$ ,  $R^{9b}$ ,  $R^{9c}$ ,  $R^{9d}$ , and  $R^{9e}$  are each independently hydrogen, alkyl, alkenyl, alkynyl, alkoxy, alkylthio, alkylsulfinyl, alkylsulfonyl, arylsulfonyl, alkoxycarbonyl, arylcarbonyl, alkylamino, arylalkyl, aryl, heterocyclic, heteroaromatic, absent, or a prodrug moiety, and  $R^{9d}$  and  $R^{9e}$  may be linked to form a ring;

$W$  is  $CR^{7d}R^{7e}$ ,  $NR^{7b}$  or O;

$W'$  is O or S; and

$R^{7a}$ ,  $R^{7b}$ ,  $R^{7c}$ ,  $R^{7d}$ , and  $R^{7e}$  are each independently hydrogen, alkyl, alkenyl, alkynyl, alkoxy, alkylthio, alkylsulfinyl, arylsulfonyl, alkoxycarbonyl, arylcarbonyl, alkylamino, arylalkyl, aryl, heterocyclic, heteroaromatic, absent, or a prodrug moiety, and  $R^{7d}$  and  $R^{7e}$  may be linked to form a ring;

and pharmaceutically acceptable salts thereof, provided that  $R^9$  is not hydrogen when  $R^7$  is dialkylamino or hydrogen.

The invention also pertains, at least in part, to methods for treating a tetracycline responsive state in a subject. The method includes administering to the subject a substituted tetracycline compound of formula (I).


In another embodiment, the invention includes pharmaceutical compositions comprising a therapeutically effective amount of a substituted tetracycline compound of formula (I) and a pharmaceutically acceptable carrier.

- 4 -

In yet another embodiment, the invention pertains to a method for synthesizing 7- and/or 9- substituted tetracycline compounds. The method includes contacting a tetracycline compound with a nitrating agent, under conditions such that a nitro tetracycline compound is formed, contacting the nitro tetracycline compound with a hydrogenating agent, under conditions such that an amino tetracycline compound is formed, and contacting the amino tetracycline compound with an amino reactive substrate, such that a 9- or 7- substituted tetracycline compound is formed.

The invention also pertains, at least in part, to a method for synthesizing a 7- and/or 9- substituted tetracycline compound of formula (I), by contacting a reactive intermediate with appropriate reagents under appropriate conditions, such that a substituted tetracycline compound of formula (I) is formed.

The reactive intermediate, wherein said reactive intermediate is of the formula:



(I)

wherein:

X is  $\text{CHC}(\text{R}^{13}\text{Y}'\text{Y})$ ,  $\text{CHR}^6$ , S,  $\text{NR}^6$ , or O;

$\text{R}^2$  is hydrogen, alkyl, alkynyl, alkoxy, alkylthio, alkylsulfinyl, alkylsulfonyl, alkylamino, arylalkyl, aryl, heterocyclic, heteroaromatic or a prodrug moiety;

$\text{R}^4$  and  $\text{R}^{4'}$  are each hydrogen, alkyl, alkenyl, alkynyl, alkoxy, alkylthio, alkylsulfinyl, alkylsulfonyl, alkylamino, arylalkyl, aryl, heterocyclic, heteroaromatic or a prodrug moiety;

$\text{R}^{2'}$ ,  $\text{R}^3$ ,  $\text{R}^{10}$ ,  $\text{R}^{11}$  and  $\text{R}^{12}$  are each hydrogen or a pro-drug moiety;

$\text{R}^5$  is hydrogen, hydroxyl, or a prodrug moiety;

$\text{R}^6$  and  $\text{R}^8$  are each independently hydrogen, alkyl, alkenyl, alkynyl, aryl, alkoxy, alkylthio, alkylsulfinyl, alkylsulfonyl, alkylamino, or an arylalkyl;

$\text{R}^{13}$  is hydrogen, hydroxy, alkyl, alkenyl, alkynyl, alkoxy, alkylthio, alkylsulfinyl, alkylsulfonyl, alkylamino; or an arylalkyl;

- 5 -

Y' and Y are each independently hydrogen; halogen; hydroxyl; cyano, sulfhydryl; amino; alkyl; alkenyl; alkynyl; alkoxy; alkylthio; alkylsulfinyl; alkylsulfonyl; alkylamino; or an arylalkyl;


R<sup>7</sup> is hydrogen, dialkylamino, thiourea, diazonium salt, thiocarboxamide, or nitro;

R<sup>9</sup> is hydrogen, thiourea, diazonium salt, thiocarboxamide, or nitro; and pharmaceutically acceptable salts thereof, provided that R<sup>9</sup> is not hydrogen when R<sup>7</sup> is hydrogen or dialkylamino.

## 10 Detailed Description Of The Invention

The present invention pertains, at least in part, to novel 7- and 9- substituted tetracycline urea, thiourea, carbamate, thiocarbamate, amino-thiazolyl, and amino-heteroaryl compounds. These compounds can be used to treat numerous tetracycline compound-responsive states, such as bacterial infections and neoplasms, as well as other known applications for tetracycline compounds, such as tetracycline efflux blockers and gene expression modulation.

In one embodiment, the invention includes 7- and 9- substituted tetracycline compounds. Preferably, the substituted tetracycline compounds are of formula (I):



20

(I)

wherein:

X is CHC(R<sup>13</sup>Y'Y), CR<sup>6</sup>R<sup>6</sup>, S, NR<sup>6</sup>, or O;

R<sup>2</sup> is hydrogen, alkyl, alkynyl, alkoxy, alkylthio, alkylsulfinyl, alkylsulfonyl, alkylamino, arylalkyl, aryl, heterocyclic, heteroaromatic or a prodrug moiety;

25

R<sup>4</sup> and R<sup>4</sup>' are each hydrogen, alkyl, alkenyl, alkynyl, alkoxy, alkylthio, alkylsulfinyl, alkylsulfonyl, alkylamino, arylalkyl, aryl, heterocyclic, heteroaromatic or a prodrug moiety;

- 6 -








- $R^{2'}$ ,  $R^3$ ,  $R^{10}$ ,  $R^{11}$  and  $R^{12}$  are each hydrogen or a pro-drug moiety;  
 $R^5$  is hydrogen, hydroxyl, or a prodrug moiety;  
 $R^6$ ,  $R^{6'}$ , and  $R^8$  are each independently hydrogen, alkyl, alkenyl, alkynyl, aryl, alkoxy, alkylthio, alkylsulfinyl, alkylsulfonyl, alkylamino, arylalkyl, or halogen;  
5  $R^7$  is hydrogen, dialkylamino, heteroaryl-amino, or  $NR^{7c}C(=W')WR^{7a}$ ;  
 $R^{13}$  is hydrogen, hydroxy, alkyl; alkenyl; alkynyl; alkoxy; alkylthio; alkylsulfinyl; alkylsulfonyl; alkylamino; or an arylalkyl;  
 $Y'$  and  $Y$  are each independently hydrogen, halogen, hydroxyl, cyano, sulfhydryl, amino, alkyl, alkenyl, alkynyl, alkoxy, alkylthio, alkylsulfinyl, alkylsulfonyl,  
10 alkylamino, or arylalkyl;  
 $R^9$  is hydrogen, heteroaryl-amino, or  $NR^{9c}C(=Z')ZR^{9a}$ ;  
 $Z$  is  $CR^{9d}R^{9e}$ ,  $NR^{9b}$ , or O;  
 $Z'$  is O or S;  
 $R^{9a}$ ,  $R^{9b}$ ,  $R^{9c}$ ,  $R^{9d}$ , and  $R^{9e}$  are each independently hydrogen, alkyl,  
15 alkenyl, alkynyl, alkoxy, alkylthio, alkylsulfinyl, alkylsulfonyl, arylsulfonyl, alkoxycarbonyl, arylcarbonyl, alkylamino, arylalkyl, aryl, heterocyclic, heteroaromatic, absent, or a prodrug moiety, and  $R^{9d}$  and  $R^{9e}$  may be linked to form a ring;  
 $W$  is  $CR^{7d}R^{7e}$ ,  $NR^{7b}$  or O;  
 $W'$  is O or S; and  
20  $R^{7a}$ ,  $R^{7b}$ ,  $R^{7c}$ ,  $R^{7d}$ , and  $R^{7e}$  are each independently hydrogen, alkyl, alkenyl, alkynyl, alkoxy, alkylthio, alkylsulfinyl, arylsulfonyl, alkoxycarbonyl, arylcarbonyl, alkylamino, arylalkyl, aryl, heterocyclic, heteroaromatic, absent, or a prodrug moiety, and  $R^{7d}$  and  $R^{7e}$  may be linked to form a ring, and pharmaceutically acceptable salts thereof, provided that  $R^9$  is not hydrogen when  $R^7$  is dialkylamino or  
25 hydrogen.

The term "tetracycline compound" includes many compounds with a similar ring structure to tetracycline. Examples of tetracycline compounds include: tetracycline, chlortetracycline, oxytetracycline, demeclocycline, methacycline, sancycline, doxycycline, and minocycline. Other derivatives and analogues comprising a similar  
30 four ring structure are also included. Table 1 depicts tetracycline and several known tetracycline derivatives. In an embodiment, the term "unsubstituted tetracycline

- 7 -

compounds," includes tetracycline compounds wherein  $R^7$  is not  $NR^{7c}C(=W')WR^{7a}$  nor heteroaryl-amino and wherein  $R^9$  is not heteroaryl-amino nor  $NR^{9c}C(=Z')ZR^{9a}$ .

TABLE I

 <p style="text-align: center;">Tetracycline</p>		
 <p style="text-align: center;">Oxytetracycline</p>	 <p style="text-align: center;">Demeclocycline</p>	 <p style="text-align: center;">Methacycline</p>
 <p style="text-align: center;">Doxycycline</p>	 <p style="text-align: center;">Chlorotetracycline</p>	 <p style="text-align: center;">Minocycline</p>

5

The term "substituted tetracycline compounds" includes tetracycline compounds with substitution at the 7- or 9- position. In one embodiment, the substitution at the 7- or 9- position enhances the ability of the substituted tetracycline compound to perform its intended function. In an embodiment, the 9- substituted tetracycline compound is 9- substituted minocycline (*e.g.*, wherein  $R^4$  and  $R^{4'}$  are methyl,  $R^5$  is hydrogen,  $R^7$  is dimethyl amino, and  $X$  is  $CR^6R^{6'}$ , wherein both  $R^6$  and  $R^{6'}$  are hydrogen atoms); 7- or 9- substituted doxycycline (*e.g.*, wherein  $R^4$  and  $R^{4'}$  are methyl,  $R^5$  is hydroxyl,  $X$  is  $CR^6R^{6'}$ ,  $R^6$  is methyl and  $R^{6'}$  is hydrogen); or a 7- or 9- substituted sancycline (wherein  $R^4$  and  $R^{4'}$  are methyl;  $R^5$  is hydrogen,  $X$  is  $CR^6R^{6'}$ ,  $R^6$  and  $R^{6'}$  are hydrogen atoms). In a further embodiment,  $R^5$  may be a protected hydroxyl group, *e.g.*, a prodrug moiety. Examples of prodrug moieties include, for example, acyl esters and propionic acid esters. In certain embodiments, the prodrug moiety is aroyl, alkanoyl, or alkaroyl and may or may not be cleaved *in vivo* to the hydroxyl group. In an embodiment,  $R^{2'}$ ,  $R^3$ ,  $R^8$ ,  $R^{10}$ ,  $R^{11}$ , and  $R^{12}$  are each hydrogen. In certain

10

15




- 8 -

embodiments of the invention, the term substituted tetracycline compounds includes tetracycline compounds wherein least one of  $R^7$  or  $R^9$  is heteroaryl-amino,  $NR^{7c}C(=W')WR^{7a}$ , or  $NR^{9c}C(=Z')ZR^{9a}$ .

The term "9-substituted tetracycline compounds" includes, in one embodiment, compounds wherein  $R^9$  is amino-heteroaryl or  $NR^{9c}C(=Z')ZR^{9a}$ . In a further  
 5 embodiment,  $R^{9c}$  is hydrogen. In another,  $Z'$  is oxygen or sulfur. In an embodiment,  $Z$  is oxygen or  $NR^{9b}$ , wherein  $R^{9b}$  is hydrogen. In another further embodiment,  $R^{9a}$  may be hydrophobic.  $R^{9a}$  may also be alkyl, alkenyl (*e.g.*, ethenyl, propenyl, butenyl, *etc.*), alkynyl, aryl (*e.g.*, phenyl, heteroaryl, *etc.*), arylalkyl, or multicyclic (*e.g.*, polycyclic,  
 10 *e.g.*, steroidyl, *e.g.*, cholesteroïdyl).

In one embodiment,  $R^{9a}$  is substituted or unsubstituted alkyl (*e.g.*, methyl, ethyl, t-butyl, n-butyl, i-butyl, or n-pentyl.) Examples of possible substituents include but are not limited to, alkyl, alkenyl, halogen, hydroxyl, alkoxy, alkylcarbonyloxy, alkyloxycarbonyl, arylcarbonyloxy, alkoxycarbonylamino, alkoxycarbonyloxy,  
 15 aryloxycarbonyloxy, carboxylate, alkylcarbonyl, alkylaminoacarbonyl, arylalkyl aminocarbonyl, alkenylaminocarbonyl, alkylcarbonyl, arylcarbonyl, aminoalkyl, arylalkylcarbonyl, alkenylcarbonyl, alkoxycarbonyl, silyl, aminocarbonyl, alkylthiocarbonyl, phosphate, aralkyl, phosphonato, phosphinato, cyano, amino, acylamino, amido, imino, sulfhydryl, alkylthio, sulfate, arylthio, thiocarboxylate,  
 20 alkylsulfinyl, sulfonato, sulfamoyl, sulfonamido, nitro, cyano, azido, heterocyclyl, alkylaryl, aryl and heteroaryl. In certain embodiments, the substituents are alkoxycarbonyl, amino, arylcarbonyl, halogen, hydroxy, alkylamino, alkoxy, or aryl. In certain embodiments, the substituent is halogen (*e.g.*, bromine, chlorine, iodine, fluorine).

25 In a further embodiment,  $R^{9a}$  includes at least one aryl group, *e.g.*, heteroaryl, phenyl, naphthyl, fluorene, *etc.* Fluorene is a moiety of the formula:



In one embodiment,  $R^{9a}$  is aryl, *e.g.*, substituted or unsubstituted phenyl. Examples of substituents include, but are not limited to, alkyl (*e.g.*, unsubstituted, *e.g.*,

- 9 -

methyl, ethyl, propyl, butyl, or substituted, *e.g.*, chloromethyl, dichloromethyl, perchloromethyl, fluoromethyl, difluoromethyl, perfluoromethyl, *etc.*), alkenyl, alkynyl, aryl, alkoxy (*e.g.*, methoxy, ethoxy, propoxy, *etc.*), aryloxy, alkylcarbonyl, arylcarbonyl, alkoxycarbonyl, aryloxycarbonyl, amido, halogen, nitro, azo, alkyl sulfonyl, and

5 arylsulfonyl.

In another embodiment, R<sup>9</sup> is amino-heteroaryl (*e.g.*, -NH-heteroaryl, *e.g.*, amino-thiazolyl). The thiazolyl substituent may be substituted with substituents such as phenyl rings. Scheme 3 below shows some representative substituents of the thiazole ring. For example, in one embodiment, the thiazole can be substituted with a

10 phenyl group, a biphenyl group, an adamantyl group, *etc.* The substituents of the thiazole can also be further substituted, *e.g.*, with an electron donating group, or an electron withdrawing group. Examples of electron withdrawing substituents include aryl groups (*e.g.*, phenyl), halogens (*e.g.*, chlorine, bromine, or fluorine), alkoxy groups (*e.g.*, methoxy, ethoxy), amines (*e.g.*, secondary amines, such as, diethylamine,

15 dimethylamine unsubstituted amines, *etc.*), nitro groups, *etc.* In other embodiments, the thiazolyl ring is linked to the tetracycline compound through an ether (-O-), alkyl, or other linkage which allows the substituted tetracycline compound to perform its intended function.

For example, in one embodiment, R<sup>9</sup> is substituted or unsubstituted heteroaryl-

20 amino, *e.g.*, thiazolyl amino. Examples of substituents of the heteroaryl include, but are not limited to, alkyl, alkenyl, halogen, hydroxyl, alkoxy, alkylcarbonyloxy, alkyloxycarbonyl, arylcarbonyloxy, alkoxycarbonylamino, alkoxycarbonyloxy, aryloxycarbonyloxy, carboxylate, alkylcarbonyl, alkylaminoacarbonyl, arylalkyl aminocarbonyl, alkenylaminocarbonyl, alkylcarbonyl, arylcarbonyl, aminoalkyl,

25 arylalkylcarbonyl, alkenylcarbonyl, alkoxycarbonyl, silyl, aminocarbonyl, alkylthiocarbonyl, phosphate, aralkyl, phosphonato, phosphinato, cyano, amino, acylamino, amido, imino, sulfhydryl, alkylthio, sulfate, arylthio, thiocarboxylate, alkylsulfinyl, sulfonato, sulfamoyl, sulfonamido, nitro, cyano, azido, heterocyclyl, alkylaryl, aryl and heteroaryl.

30 In certain embodiments, the substituents of the thiazolyl include alkyl (*e.g.*, methyl, ethyl, propyl, butyl, *etc.*), halogen (*e.g.*, fluorine, bromine, chlorine, iodine,

- 10 -

*etc.*), alkoxy (*e.g.*, methoxy, propoxy, ethoxy, *etc.*), aryl (*e.g.*, substituted or unsubstituted phenyl or heteroaryl).

In an embodiment, the aryl thiazolyl substituent is substituted with one or more substituents. Examples of substituents include, but are not limited to, alkyl (*e.g.*,  
 5 methyl, ethyl, propyl, butyl, *etc.*), alkenyl, alkynyl, aryl, alkoxy, aryloxy, alkylcarbonyl, arylcarbonyl, amido, trifluoromethyl, halogen, nitro, azo, alkyl sulfonyl, alkoxycarbonyl (*e.g.*, methoxycarbonyl, ethoxycarbonyl, *etc.*), aryloxycarbonyl, and arylsulfonyl.

The term "7-substituted tetracycline compounds" includes tetracycline compounds with substitution at the 7 position. Examples of tetracycline compounds  
 10 which advantageously may be substituted at the 7 position include tetracycline, sancycline, doxycycline, oxytetracycline, demeclocycline, or methacycline. In an advantageous embodiment,  $R^7$  is  $NR^{7c}C(=W')WR^{7a}$ , wherein  $R^{7c}$  may be hydrogen,  $W'$  may be oxygen or sulfur.  $R^{7a}$  may be hydrophobic.  $R^{7a}$  may also be alkyl, alkenyl (*e.g.*, ethenyl, propenyl, butenyl, *etc.*), alkynyl, aryl (*e.g.*, phenyl, heteroaryl, *etc.*), arylalkyl,  
 15 heteroaromatic, or multicyclic (*e.g.*, polycyclic, *e.g.*, steroidyl, *e.g.*, cholesteroildyl).  $R^{7a}$  may also include at least one phenyl group, *e.g.*, naphthyl or fluorene. In a preferred embodiment,  $W$  is oxygen or  $NR^{7b}$ , wherein  $R^{7b}$  is hydrogen.

In one embodiment,  $R^{7a}$  is substituted or unsubstituted alkyl (*e.g.*, methyl, ethyl, t-butyl, n-butyl, i-butyl, or n-pentyl.) Examples of possible substituents include but are  
 20 not limited to, alkyl, alkenyl, halogen, hydroxyl, alkoxy, alkylcarbonyloxy, alkylloxycarbonyl, arylcarbonyloxy, alkoxycarbonylamino, alkoxycarbonyloxy, aryloxycarbonyloxy, carboxylate, alkylcarbonyl, alkylaminoacarbonyl, arylalkyl aminocarbonyl, alkenylaminocarbonyl, alkylcarbonyl, arylcarbonyl, aminoalkyl, arylalkylcarbonyl, alkenylcarbonyl, alkoxycarbonyl, silyl, aminocarbonyl,  
 25 alkylthiocarbonyl, phosphate, aralkyl, phosphonato, phosphinato, cyano, amino, acylamino, amido, imino, sulfhydryl, alkylthio, sulfate, arylthio, thiocarboxylate, alkylsulfinyl, sulfonato, sulfamoyl, sulfonamido, nitro, cyano, azido, heterocyclyl, alkylaryl, aryl and heteroaryl. In certain embodiments, the substituents are alkoxycarbonyl, amino, arylcarbonyl, halogen, hydroxy, alkylamino, alkoxy, or aryl. In  
 30 certain embodiments, the substituent is halogen (*e.g.*, bromine, chlorine, iodine, fluorine). In a further embodiment,  $R^{7a}$  includes at least one aryl group, *e.g.*, heteroaryl, phenyl, naphthyl, fluorene, *etc.*

- 11 -

In one embodiment, R<sup>7a</sup> is aryl, *e.g.*, substituted or unsubstituted phenyl. Examples of substituents include, but are not limited to, alkyl (*e.g.*, unsubstituted, *e.g.*, methyl, ethyl, propyl, butyl, or substituted, *e.g.*, chloromethyl, dichloromethyl, perchloromethyl, fluoromethyl, difluoromethyl, perfluoromethyl, *etc.*), alkenyl, alkynyl, 5 aryl, alkoxy (*e.g.*, methoxy, ethoxy, propoxy, *etc.*), aryloxy, alkylcarbonyl, arylcarbonyl, alkoxycarbonyl, aryloxycarbonyl, amido, halogen, nitro, azo, alkyl sulfonyl, and arylsulfonyl.

In another embodiment, R<sup>7</sup> is amino-heteroaryl (*e.g.*, -NH-heteroaryl, *e.g.*, amino-thiazolyl). The thiazolyl substituent may be substituted with substituents such 10 as phenyl rings. Scheme 3 below shows some representative substituents of the thiazole ring. For example, in one embodiment, the thiazole can be substituted with a phenyl group, a biphenyl group, an adamantyl group, *etc.* The substituents of the thiazole can also be further substituted, *e.g.*, with an electron donating group, or an electron withdrawing group. Examples of electron withdrawing substituents include 15 aryl groups (*e.g.*, phenyl), halogens (*e.g.*, chlorine, bromine, or fluorine), alkoxy groups (*e.g.*, methoxy, ethoxy), amines (*e.g.*, secondary amines, such as, diethylamine, dimethylamine unsubstituted amines, *etc.*), nitro groups, *etc.* In other embodiments, the thiazolyl ring is linked to the tetracycline compound through an ether (-O-), alkyl, or other linkage which allows the substituted tetracycline compound to perform its 20 intended function.

For example, in one embodiment, R<sup>7</sup> is substituted or unsubstituted heteroaryl-amino, *e.g.*, thiazolyl amino. Examples of substituents of the heteroaryl include, but are not limited to, alkyl, alkenyl, halogen, hydroxyl, alkoxy, alkylcarbonyloxy, alkylloxycarbonyl, arylcarbonyloxy, alkoxycarbonylamino, alkoxycarbonyloxy, 25 aryloxycarbonyloxy, carboxylate, alkylcarbonyl, alkylaminoacarbonyl, arylalkyl aminocarbonyl, alkenylaminocarbonyl, alkylcarbonyl, arylcarbonyl, aminoalkyl, arylalkylcarbonyl, alkenylcarbonyl, alkoxycarbonyl, silyl, aminocarbonyl, alkylthiocarbonyl, phosphate, aralkyl, phosphonato, phosphinato, cyano, amino, acylamino, amido, imino, sulfhydryl, alkylthio, sulfate, arylthio, thiocarboxylate, 30 alkylsulfinyl, sulfonato, sulfamoyl, sulfonamido, nitro, cyano, azido, heterocyclyl, alkylaryl, aryl and heteroaryl.

- 12 -

In certain embodiments, the substituents of the thiazolyl include alkyl (*e.g.*, methyl, ethyl, propyl, butyl, *etc.*), halogen (*e.g.*, fluorine, bromine, chlorine, iodine, *etc.*), alkoxy (*e.g.*, methoxy, propoxy, ethoxy, *etc.*), aryl (*e.g.*, substituted or unsubstituted phenyl or heteroaryl).

5           In an embodiment, the aryl thiazolyl substituent is substituted with one or more substituents. Examples of substituents include, but are not limited to, alkyl (*e.g.*, methyl, ethyl, propyl, butyl, *etc.*), alkenyl, alkynyl, aryl, alkoxy, aryloxy, alkylcarbonyl, arylcarbonyl, amido, trifluoromethyl, halogen, nitro, azo, alkyl sulfonyl, alkoxycarbonyl (*e.g.*, methoxycarbonyl, ethoxycarbonyl, *etc.*), aryloxycarbonyl, and arylsulfonyl.

10           In a further embodiment of the invention,  $R^2$ ,  $R^3$ ,  $R^{10}$ ,  $R^{11}$ , and  $R^{12}$  of the substituted tetracycline compounds are each hydrogen atoms. In another further embodiment,  $R^4$  and  $R^4$  are each alkyl, *e.g.*, lower alkyl, and, advantageously, methyl. In another embodiment, X is  $CR^6R^{6'}$ .  $R^6$  and  $R^{6'}$  are selected from the group consisting of hydrogen, methyl, and hydroxy groups.

15           In another embodiment, the invention also pertains to compounds wherein both  $R^9$  and  $R^7$  are not hydrogen. These compounds may be referred to as 7- and 9-disubstituted compounds. The invention pertains to compounds with any combination of 7- and 9- substituents disclosed herein.

20           Examples of compounds of the invention include those disclosed in Table 2, in addition to the compounds listed below. The invention also pertains to pharmaceutically acceptable salts of any of these compounds as well as enantiomers, and mixtures of the compounds. Examples of compounds of the invention include, but are not limited to:

25           Doxycycline 9-carbamic acid 9*H*-fluoren-9-ylmethyl ester;  
          (9-(Naphthyn-1-yl) doxycycline urea;  
          9-(3-Methyl-1-butyl) doxycycline urea;  
          9-Phenyl doxycycline urea;  
          9-*t*-Butyl doxycycline urea;  
          Fmoc 9-amino doxycycline;  
          9-(4'-Chloro-2'-trifluoromethylphenyl) doxycycline urea;  
30           9-(4'-Fluorophenyl) doxycycline carbamate;  
          9-(4'-Methoxyphenyl) doxycycline carbamate;  
          9-BOC amino doxycycline;

- 13 -

- 9-(Phenylthiazolyl) amino doxycycline;  
9-(Ethylthiazolyl) amino doxycycline;  
(4-Fluorophenylthiazolyl) amino doxycycline;  
9-(4'-Methoxyphenylthiazolyl) amino doxycycline;  
5 9-(3'-Nitrophenylthiazolyl) amino doxycycline;  
9-(4'-Methyl, 5'-phenylthiazolyl) amino doxycycline;  
9-Neopentyl minocycline carbamate;  
9-(Phenylthiazolyl) amino sancycline;  
9-(Adamantylthiazolyl) amino doxycycline;  
10 9-(Naphthyn-1-yl urea) Doxycycline 5-propanoic acid ester;  
Doxycycline 9-Thiocarbamic acid 9*H*-fluoren-9-ylmethyl ester;  
(9-(Naphthyn-1-yl) doxycycline thiourea;  
9-(3-methyl-1-butyl) doxycycline thiourea;  
9-Phenyl doxycycline thiourea;  
15 9-*t*-Butyl doxycycline thiourea;  
9-(4'-Chloro-2'-trifluoromethylphenyl) doxycycline thiourea;  
9-(4'-Fluorophenyl) doxycycline thiocarbamate;  
9-(4-Methoxyphenyl) doxycycline thiocarbamate;  
9-Neopentyl minocycline thiocarbamate;  
20 9-(Naphthyn-1-yl) doxycycline thiourea 5-propanoic acid ester;  
Minocycline 9-carbamic acid 9*H*-fluoren-9-ylmethyl ester;  
(9-(Naphthyn-1-yl) minocycline urea;  
9-(3-Methyl-1-butyl) minocycline urea;  
9-Phenyl doxycycline urea;  
25 9-*t*-Butyl minocycline urea;  
FMOC 9-amino minocycline;  
9-(4'-Chloro-2'-trifluoromethylphenyl) minocycline urea;  
9-(4'-Fluorophenyl) minocycline carbamate;  
9-(4'-Methoxyphenyl) minocycline carbamate;  
30 9-BOC amino minocycline;  
9-(Phenylthiazolyl) amino minocycline;  
9-(Ethylthiazolyl) amino minocycline;

- 14 -

- (4'-Fluorophenylthiazolyl) amino minocycline;  
9-(4'-Methoxyphenylthiazolyl) amino minocycline;  
9-9-(3'-Nitrophenylthiazolyl) amino minocycline;  
9-(4'-Methyl, 5'-phenylthiazolyl) amino doxycycline;  
5 9-Neopentyl doxycycline carbamate;  
9-(Phenylthiazolyl) amino minocycline;  
9-(Adamantylthiazolyl) amino minocycline;  
Minocycline 9-thiocarbamic acid 9*H*-fluoren-9-ylmethyl ester;  
(9-(Naphthyn-1-yl) minocycline thiourea;  
10 9-(3'-Methyl-1-butyl) minocycline thiourea;  
9-Phenyl minocycline thiourea;  
9-*t*-Butyl minocycline thiourea;  
9-(4'-Fluorophenyl) minocycline thiocarbamate;  
9-(4'-Methoxyphenyl) minocycline thiocarbamate;  
15 9-Neopentyl doxycycline thiocarbamate;  
9-(2'-Bromoethyl) doxycycline carbamate;  
9-(*n*-Pentyl) minocycline carbamate;  
9-(4'-Benzoylbenzoyl) amino doxycycline;  
7-(3'-Nitrophenylthiazolyl) amino sancycline;  
20 9-(3'-Ethoxycarbonylthiazolyl) amino doxycycline;  
7-(4'-Methylphenyl) sancycline carbamate;  
9-(4'-Trifluoromethoxyphenyl) minocycline urea;  
9-(3', 5'-diperfluorophenyl) minocycline thiourea;  
9-Prop-2'-enyl minocycline carbamate;  
25 9-(4'-Chloro, 2'-nitrophenyl) minocycline urea;  
9-Ethyl minocycline carbamate;  
9-*n*-Butyl minocycline carbamate;  
9-*n*-But-3-enyl minocycline carbamate;  
Doxycycline 7-carbamic acid 7*H*-fluoren-7-ylmethyl ester;  
30 7-(Naphthyn-1-yl) doxycycline urea;  
7-(3-Methyl-1-butyl) doxycycline urea;  
7-Phenyl doxycycline urea;



















- 15 -

- 7-t-Butyl doxycycline urea;  
7-Fmoc amino doxycycline;  
7-(4'-Chloro-2-trifluoromethylphenyl) doxycycline urea;  
7-(4'-Fluorophenyl) doxycycline carbamate;  
5 7-(4'-Methoxyphenyl) doxycycline carbamate;  
7-BOC amino doxycycline;  
7-(3'-Phenylthiazolyl) amino doxycycline;  
7-(3'-Ethylthiazolyl) amino doxycycline;  
7-(4''-Fluorophenylthiazolyl) amino doxycycline;  
10 7-(4''-Methoxyphenylthiazolyl) amino doxycycline;  
7-(Phenylthiazolylamino)-sancycline;  
7-(3'-Nitrophenylthiazolyl) amino doxycycline;  
7-(4'-Methyl, 5'-phenylthiazolyl) amino doxycycline;  
7-(Adamantylthiazolyl) amino doxycycline;  
15 Doxycycline 7-thiocarbamic acid 7*H*-fluoren-7-ylmethyl ester;  
7-(Naphthyn-1-yl) doxycycline thiourea;  
7-(3-Methyl-1-butyl) doxycycline thiourea;  
7-Phenyl amino doxycycline thiourea;  
7-t-butyl amino doxycycline thiourea;  
20 7-(4'-Chloro-2'-trifluoromethylphenyl) doxycycline thiourea;  
7-(4'-Fluorophenyl) doxycycline thiocarbamate;  
7-(4'-Methoxyphenyl) doxycycline thiocarbamate;  
7-(Naphthyn-1-yl) doxycycline urea 5-propanoic acid ester;  
9-i-Butyl minocycline carbamate,  
25 7-(Naphthyn-1-yl) doxycycline thiourea 5-propanoic acid ester, and  
pharmaceutically acceptable salts thereof.













Other compounds of the invention include compounds having the following structures:




- 16 -



- 17 -



- 18 -



The invention includes methods for synthesizing 7- and/or 9- substituted tetracycline compounds using reactive intermediates, such as thioureas, thiocarboxyamides, and diazonium salts, advantageously, at the R<sup>7</sup> and/or R<sup>9</sup> position of tetracycline compound of formula (I). In one embodiment, the invention pertains to methods of synthesizing 9- and/or 7- substituted tetracycline compounds by contacting a reactive intermediate with appropriate reagents under appropriate conditions, such that a 7- and/or 9- substituted tetracycline compound is formed. Examples of appropriate reagents and conditions are described in Schemes 1-3 and in Example 1.

The term "reactive intermediates" includes species which are generated during the synthesis of the 7- and/or 9- substituted tetracycline compounds of Formula (I). These intermediates may or may not be stable under reaction conditions and may or may not be isolatable. However, one of skill in the art can appreciate that these reactive intermediates can be used to generate other 7- and/or 9- substituted tetracycline compounds. The term "reactive intermediates" includes all intermediates which are synthesized or can be synthesized using the methodology discussed herein. Examples of reactive intermediates of the invention include 7- and/or 9- diazonium salts, 7- and/or 9- thiocarboxyamates, 7- and/or 9- anilino compounds, 7- and/or 9- amino tetracycline compounds, 7- and/or 9- nitro tetracycline compounds, 7- and/or 9- urea derivatives, 7- and/or 9- carbamate derivatives, *etc.*

The language "appropriate conditions" include conditions known in the art and conditions described herein to convert the reactive intermediate to a 7- and/or 9- substituted tetracycline compound of formula (I) or another desired tetracycline compound.

The language "appropriate reagents" include reagents known in the art and reagents described herein to convert the reactive intermediate to a tetracycline compound of formula (I) or another desired tetracycline compound.

- 19 -

In another embodiment, the invention includes methods of synthesizing 7- and/or 9- substituted tetracycline compounds outlined in the following schemes. Although in each scheme the reaction is shown for only one or two tetracycline compounds, one of skill in the art will appreciate that similar reactions can be also be performed with other  
5 tetracycline compounds.

In an embodiment, the invention pertains to a method for synthesizing 7- and/or 9-substituted tetracycline compounds by contacting an unsubstituted tetracycline compound with a nitrating agent, to form a 7- and/or 9-nitro substituted tetracycline compound. The 7- and/or 9-nitro substituted tetracycline compound is then  
10 hydrogenated with a hydrogenating agent to form a 7- and/or 9-amino substituted tetracycline compound. The 7- and/or 9-amino substituted tetracycline compound is contacted with an amino reactive compound, thereby forming a 7- and/or 9-substituted tetracycline compound.


The term "nitrating agent" includes compounds and chemicals which, under  
15 appropriate conditions, can introduce a nitro group ( $-\text{NO}_2$ ) to a tetracycline compound. Advantageous nitrating agents include, for example,  $\text{NaNO}_2$ . Other methods of nitration are known in the art and are also included (see, for example, March, *Advanced Organic Chemistry*, John Wiley & Sons:New York, 1992, p. 522-525, and references cited therein). Advantageously, when the nitrating agent is  $\text{NaNO}_2$ , the reaction is conducted  
20 under acidic conditions.

The term "hydrogenating agent" includes compounds and chemicals which, under appropriate conditions, can convert a nitro group to an amino group ( $-\text{NH}_2$ ). Examples of preferred hydrogenating agents include  $\text{H}_2$  gas with a transition metal catalyst, advantageously, platinum. Other methods of converting nitro groups to amino  
25 groups are known in the art (see, for example March, *Advanced Organic Chemistry*, John Wiley & Sons:New York, 1992, p. 1216-1217, and references cited therein).

The term "amino reactive compound" includes compounds and molecules which can be reacted with the 7-and/or 9-amino tetracycline derivative to form a desired 7- and/or 9-substituted tetracycline compound, or a precursor thereof. Examples of  
30 advantageous amino reactive compounds for the formation of 9-substituted urea and carbamate tetracycline compounds include substituted and unsubstituted isocyanates and chloroformates. Scheme 1 below depicts the synthesis for a 9-substituted doxycycline

compounds, but the methodology can be applied both to other tetracycline compounds and 7- substituted tetracycline compound. The depicted method includes treating an unsubstituted tetracycline compound (1-1) with acid (*e.g.*,  $\text{H}_2\text{SO}_4$ ) and a nitrating agent (*e.g.*, sodium or potassium nitrate) to form the reactive nitro substituted tetracycline intermediate (1-2). The reactive nitro substituted tetracycline intermediate can be reduced to the corresponding amine (1-3) by hydrogenating reagents known in the art (*e.g.*, hydrogen with metal catalysts, platinum oxide or the like) to produce the amino substituted tetracycline compound (1-3). The amino substituted tetracycline compound (1-3) can then be reacted in mild base with substituted isocyanates (1-4) to form mixed urea substituted tetracycline compounds (1-5). The amino substituted tetracycline compounds (1-3) can also be reacted with substituted or unsubstituted chloroformates (1-6) to form substituted carbamates tetracycline compounds (1-7). Additionally, the amino substituted tetracycline compounds (1-3) can be reacted with other species, such as the thioisocyanates (1-8), to form other desirable derivatives, such as the thioureas (1-9).

SCHEME 1:



- 21 -


The initial nitration of the tetracycline compound may produce a mixture of the 7- and 9-substituted isomers. An ordinarily skilled artisan will be able to appreciate that the isomers can be separated by conventional methods after any of the reactions mentioned above. Techniques for separating isomers are well known in the art. For example, the amino substituted tetracycline compounds (*e.g.*, 1-3) can be separated from other positional isomers by techniques known in the art, *e.g.*, preparative HPLC on C18-reverse phase silica gel with a binary gradient system. The amino tetracycline compounds can also be prepared according to U.S. Patent No. 3,483,251 through a reductive alkylation of 7-(N,N''-dicarbobenzyloxyhydrazino) tetracyclines.

Furthermore, other 7- and/or 9-substituted tetracycline compounds can be synthesized by reacting the amino intermediate with amino reactive substrate.

The reactive 7- and/or 9- amino substituted tetracycline compounds are included as reactive intermediates. The amino substituted tetracycline compounds can react with other chemical species such as isocyanate derivatives or isothiocyanate derivatives to produce 7- and/or 9- position ureas and thioureas (thiocarbocarboxyamides) as shown in Scheme 1. The 7- and/or 9- position urea and thiourea tetracycline compounds are reactive intermediates and can be used in the synthesis of a wide variety of 7- and/or 9-substituted tetracycline compounds. For example, the 7- and/or 9- position thioureas can be used to form amino-heterocyclic moieties by reactions shown in Scheme 2, below.

- 22 -

## SCHEME 2:




- For example, 9-amino substituted tetracycline compound (2-1) can be reacted
- 5 with Fmoc-NCS to produce a 9-Fmoc thiourea substituted tetracycline compound (2-2). The Fmoc substituted thiourea substituted tetracycline compound (2-2) can be deprotected using methods known in the art to form the 9-thiourea substituted tetracycline compound (2-3). The 9-thiourea substituted tetracycline compound (2-3) is a reactive intermediate, which can be reacted with  $\alpha$ -haloketones (2-4, *e.g.*, substituted
- 10 or unsubstituted  $\alpha$ -haloketones, *etc.*), to produce 9-thiazolylamino substituted tetracycline compounds (2-5). This methodology can also be used to form 7- position thiourea substituted tetracycline reactive intermediates as well as 7- position thiazolylamino tetracycline compounds.

- Thiourea tetracycline reactive intermediates also can be used as reactive
- 15 intermediates in the synthesis of, for example, spiro and fused cyclopentapyrrole and

- 23 -

pyrimidine derivatives (Albar *et al.*, *J. Chem. Res., Synop.*, (2), 40-41 (1997);  
 pyridazinedione derivatives (Sharaf El-Din, *Alexandria J. Pharm. Sci.*, 11(1) 9-12  
 (1997)); benzothiazole acrylic acid derivatives (Kassem, *et al.*, *Pak. J. Sci. Ind. Res.* 38  
 (11-12) 424-427 (1995)); thiazoline, arylazothioazole and pyrazole derivatives  
 5 (Abdelhamid, A. *Phosphorous, Sulfur, Silicon Related Elem.* 119 (181-191) (1996));  
 pyrimidine derivatives (Fikry, *J. Indian Chem. Soc.*, 73(12), 698-699 (1996));  
 aminothiazole-carbonitrile derivatives (Shiono, JP 95-331456); benzimidazole  
 derivatives (Omar *et al. Egypt. J. Pharm. Sci.* 37(1-6), 609-620 (1996));  
 benzylthiazolidine derivatives (Morita *et al.*, JP 95-200268); clonidine derivatives  
 10 (Pierce, *et al.* WO 95/21818); pyrimidine derivatives (Nassar, *et al. Egypt. J. Chem.*,  
 40(3) 239-247 (1997)); bicyclic derivatives (Zhu, *et al. Hanneng Cailiao* 5(4), 165-170  
 (1997)); combinatorial libraries (Nefzi, *et al.* WO 98/19693); triazinoindole derivatives  
 (Tomchin, *et al. Khim. -Farm. Zh.*, 31(3), 19-27 (1997); Tomchin *et al.*, *Khim.-Farm.*  
*Zh.*, 32(3) 7-10 (1998)); aryl thio derivatives (Chikalia, *et al. Proc. Nat. Acad. Sci. India*  
 15 68(A), I, 1998); and  $\alpha$ -amino acid derivatives (Beyer, *et al. Tetrahedron*, 52(17) 6233-  
 6240 (1996)).

## SCHEME 3:




As depicted in Scheme 3, 5-esters of 7- and/or 9- substituted tetracycline  
 20 compounds (3-1 and 3-3) can be formed by dissolving the 7- and/or 9- substituted  
 tetracycline compounds in strong acid and adding the appropriate carboxylic acid.



- 24 -

Examples of strong acids include anhydrous hydrogen fluoride, methanesulphonic acid, and trifluoromethanesulfonic acid.

The invention also pertains to a method for synthesizing a 7- or 9- substituted tetracycline compound of formula (I), by contacting a reactive intermediate with  
 5 appropriate reagents under appropriate conditions, such that a substituted tetracycline compound is formed. The compound of formula (I) is:



(I)

wherein:

- 10 X is  $\text{CHC}(\text{R}^{13}\text{Y}'\text{Y})$ ,  $\text{CR}^6\text{R}^6$ , S,  $\text{NR}^6$ , or O;  
 $\text{R}^2$  is hydrogen, alkyl, alkynyl, alkoxy, alkylthio, alkylsulfinyl, alkylsulfonyl, alkylamino, arylalkyl, aryl, heterocyclic, heteroaromatic or a prodrug moiety;  
 $\text{R}^4$  and  $\text{R}^{4'}$  are each hydrogen, alkyl, alkenyl, alkynyl, alkoxy, alkylthio,  
 15 alkylsulfinyl, alkylsulfonyl, alkylamino, arylalkyl, aryl, heterocyclic, heteroaromatic or a prodrug moiety;  
 $\text{R}^{2'}$ ,  $\text{R}^3$ ,  $\text{R}^{10}$ ,  $\text{R}^{11}$  and  $\text{R}^{12}$  are each hydrogen or a pro-drug moiety;  
 $\text{R}^5$  is hydrogen, hydroxyl, or a prodrug moiety;  
 $\text{R}^6$ ,  $\text{R}^{6'}$ , and  $\text{R}^8$  are each independently hydrogen, alkyl, alkenyl, alkynyl,  
 20 aryl, alkoxy, alkylthio, alkylsulfinyl, alkylsulfonyl, alkylamino, arylalkyl, or halogen;  
 $\text{R}^7$  is hydrogen, dialkylamino, heteroaryl-amino, or  $\text{NR}^{7c}\text{C}(=\text{W}')\text{WR}^{7a}$ ;  
 $\text{R}^{13}$  is hydrogen, hydroxy, alkyl; alkenyl; alkynyl; alkoxy; alkylthio; alkylsulfinyl; alkylsulfonyl; alkylamino; or an arylalkyl;  
 $\text{Y}'$  and  $\text{Y}$  are each independently hydrogen; halogen; hydroxyl; cyano,  
 25 sulfhydryl; amino; alkyl; alkenyl; alkynyl; alkoxy; alkylthio; alkylsulfinyl; alkylsulfonyl; alkylamino; or an arylalkyl;  
 $\text{R}^9$  is hydrogen,  $\text{NR}^{9c}\text{C}(=\text{Z}')\text{ZR}^{9a}$ , or heteroaryl-amino;  
 $\text{Z}$  is  $\text{CR}^{9d}\text{R}^{9e}$ ,  $\text{NR}^{9b}$ , or O;  
 $\text{Z}'$  is O or S;

- 25 -

$R^{9a}$ ,  $R^{9b}$ ,  $R^{9c}$ ,  $R^{9d}$ , and  $R^{9e}$  are each independently hydrogen, alkyl, alkenyl, alkynyl, alkoxy, alkylthio, alkylsulfinyl, alkylsulfonyl, arylsulfonyl, alkoxycarbonyl, arylcarbonyl, alkylamino, arylalkyl, aryl, heterocyclic, heteroaromatic, absent, or a prodrug moiety;

5 W is  $CR^{7d}R^{7e}$ ,  $NR^{7b}$  or O;

$W'$  is O or S; and


$R^{7a}$ ,  $R^{7b}$ ,  $R^{7c}$ ,  $R^{7d}$ , and  $R^{7e}$  are each independently hydrogen, alkyl, alkenyl, alkynyl, alkoxy, alkylthio, alkylsulfinyl, arylsulfonyl, alkoxycarbonyl, arylcarbonyl, alkylamino, arylalkyl, aryl, heterocyclic, heteroaromatic, absent, or a  
10 prodrug moiety;

and pharmaceutically acceptable salts thereof, provided that  $R^9$  is not hydrogen when  $R^7$  is dialkylamino or hydrogen.

In one embodiment, the reactive intermediate is a 7- and/or 9- diazonium salt, a 7- and/or 9- nitro compound, a 7- and/or 9- thiourea, or a 7- and/or 9- thiocarboxamide.

15

The invention also pertains to reactive intermediates of the formula:



(I)

wherein:

20 X is  $CHC(R^{13}Y'Y)$ ,  $CHR^6$ , S,  $NR^6$ , or O;

$R^2$  is hydrogen, alkyl, alkynyl, alkoxy, alkylthio, alkylsulfinyl, alkylsulfonyl, alkylamino, arylalkyl, aryl, heterocyclic, heteroaromatic or a prodrug moiety;

$R^4$  and  $R^{4'}$  are each hydrogen, alkyl, alkenyl, alkynyl, alkoxy, alkylthio, alkylsulfinyl, alkylsulfonyl, alkylamino, arylalkyl, aryl, heterocyclic, heteroaromatic or  
25 a prodrug moiety;

$R^{2'}$ ,  $R^3$ ,  $R^{10}$ ,  $R^{11}$  and  $R^{12}$  are each hydrogen or a pro-drug moiety;

$R^5$  is hydrogen, hydroxyl, or a prodrug moiety;

- 26 -

$R^6$  and  $R^8$  are each independently hydrogen, alkyl, alkenyl, alkynyl, aryl, alkoxy, alkylthio, alkylsulfinyl, alkylsulfonyl, alkylamino, or an arylalkyl;

$R^{13}$  is hydrogen, hydroxy, alkyl, alkenyl, alkynyl, alkoxy, alkylthio, alkylsulfinyl, alkylsulfonyl, alkylamino; or an arylalkyl;

5            $Y'$  and  $Y$  are each independently hydrogen; halogen; hydroxyl; cyano, sulfhydryl; amino; alkyl; alkenyl; alkynyl; alkoxy; alkylthio; alkylsulfinyl; alkylsulfonyl; alkylamino; or an arylalkyl;

$R^9$  is hydrogen, thiourea, diazonium salt, thiocarboxamide, or nitro;

10            $R^7$  is hydrogen, thiourea, dialkylamino, diazonium salt, thiocarboxamide, or nitro; and pharmaceutically acceptable salts thereof, provided that  $R^9$  is not hydrogen when  $R^7$  hydrogen or dialkylamino.

In a further embodiment,  $R^7$  is hydrogen or dialkylamino, when  $R^9$  is thiourea, diazonium salt, thiocarboxamide, or a nitro moiety. In another,  $R^9$  is hydrogen when  $R^7$  is thiourea, diazonium salt, thiocarboxamide, or a nitro moiety.

15           Unless specifically indicated, the chemical groups of the present invention may be substituted or unsubstituted. Further, unless specifically indicated, the chemical substituents may in turn be substituted or unsubstituted. In addition, multiple substituents may be present on a chemical group or substituent. Examples of substituents include alkyl, alkenyl, alkynyl, halogen, hydroxyl, alkylcarbonyloxy, 20 arylcarbonyloxy, alkoxy carbonyloxy, aryloxy carbonyloxy, carboxyl, alkylcarbonyl, arylcarbonyl, alkoxy carbonyl, aminocarbonyl, alkylaminocarbonyl, dialkylaminocarbonyl, alkylthiocarbonyl, alkoxy, formyl, trimethylsilyl, phosphate, phosphonate, phosphinate, cyano, amino (including alkyl amino, dialkylamino, arylamino, diarylamino, and alkylarylamino), acylamino (including alkylcarbonylamino, 25 arylcarbonylamino, carbamoyl and ureido), amido, imino, sulfhydryl, alkylthio, arylthio, thiocarboxylate, sulfates, alkylsulfinyl, sulfonate, sulfamoyl, sulfonamido, nitro, trifluoromethyl, cyano, azido, heterocyclyl, alkylaryl, aromatic or heteroaromatic moieties, cholesterol, arylsulfonyl, azo, thiazolyl, adamantyl, and phosphonyl.

30           The term "alkyl" includes saturated aliphatic groups, including straight-chain alkyl groups, branched-chain alkyl groups, cycloalkyl (alicyclic) groups, alkyl substituted cycloalkyl groups, and cycloalkyl substituted alkyl groups. The term alkyl further includes alkyl groups, which can further include oxygen, nitrogen, sulfur or

phosphorous atoms replacing one or more carbons of the hydrocarbon backbone, *e.g.*, oxygen, nitrogen, sulfur or phosphorous atoms. In an embodiment, a straight chain or branched chain alkyl has 10 or fewer carbon atoms in its backbone (*e.g.*, C<sub>1</sub>-C<sub>10</sub> for straight chain, C<sub>3</sub>-C<sub>10</sub> for branched chain), and in another embodiment, 4 or fewer.

- 5 Likewise, in certain embodiments, cycloalkyls have from 4-7 carbon atoms in their ring structure, and may have 5 or 6 carbons in the ring structure.

Moreover, the term alkyl includes both "unsubstituted alkyls" and "substituted alkyls", the latter of which refers to alkyl moieties having substituents replacing a hydrogen on one or more carbons of the hydrocarbon backbone. Such substituents can include, for example, halogen, hydroxyl, alkylcarbonyloxy, arylcarbonyloxy, 10 alkoxy carbonyloxy, aryloxy carbonyloxy, carboxylate, alkylcarbonyl, alkoxy carbonyl, aminocarbonyl, alkylthiocarbonyl, alkoxy, phosphate, phosphonate, phosphinato, cyano, amino (including alkyl amino, dialkylamino, arylamino, diarylamino, and alkylaryl amino), acylamino (including alkylcarbonylamino, arylcarbonylamino, 15 carbamoyl and ureido), amidino, imino, sulfhydryl, alkylthio, arylthio, thiocarboxylate, sulfates, sulfonate, sulfamoyl, sulfonamido, nitro, trifluoromethyl, cyano, azido, heterocyclyl, alkylaryl, or an aromatic or heteroaromatic moiety. Cycloalkyls can be further substituted, *e.g.*, with the substituents described above. An "alkylaryl" moiety is an alkyl substituted with an aryl (*e.g.*, phenylmethyl (benzyl)).

- 20 The term "aryl" includes aryl groups, including 5- and 6-membered single-ring aromatic groups that may include from zero to four heteroatoms, for example, benzene, pyrrole, furan, thiophene, imidazole, benzoxazole, benzothiazole, triazole, tetrazole, pyrazole, pyridine, pyrazine, pyridazine and pyrimidine, and the like. Aryl groups also include polycyclic fused aromatic groups such as naphthyl, quinolyl, indolyl, and the 25 like. Those aryl groups having heteroatoms in the ring structure may also be referred to as "aryl heterocycles", "heteroaryls" or "heteroaromatics". The aromatic ring can be substituted at one or more ring positions with such substituents as described above, as for example, halogen, hydroxyl, alkoxy, alkylcarbonyloxy, arylcarbonyloxy, alkoxy carbonyloxy, aryloxy carbonyloxy, carboxylate, alkylcarbonyl, alkoxy carbonyl, aminocarbonyl, alkylthiocarbonyl, phosphate, phosphonate, phosphinato, cyano, amino 30 (including alkyl amino, dialkylamino, arylamino, diarylamino, and alkylaryl amino), acylamino (including alkylcarbonylamino, arylcarbonylamino, carbamoyl and ureido),

- 28 -

amidino, imino, sulfhydryl, alkylthio, arylthio, thiocarboxylate, sulfates, sulfonato, sulfamoyl, sulfonamido, nitro, trifluoromethyl, cyano, azido, heterocyclyl, alkylaryl, or an aromatic or heteroaromatic moiety. Aryl groups can also be fused or bridged with alicyclic or heterocyclic rings which are not aromatic so as to form a polycycle (*e.g.*,  
5 tetralin).

The terms "alkenyl" and "alkynyl" include unsaturated aliphatic groups analogous in length and possible substitution to the alkyls described above, but that contain at least one double or triple bond, respectively.

Unless the number of carbons is otherwise specified, "lower alkyl" as used  
10 herein means an alkyl group, as defined above, but having from one to five carbon atoms in its backbone structure. Likewise, "lower alkenyl" and "lower alkynyl" have similar chain lengths.

The terms "alkoxyalkyl", "polyaminoalkyl" and "thioalkoxyalkyl" include alkyl groups, as described above, which further include oxygen, nitrogen or sulfur atoms  
15 replacing one or more carbons of the hydrocarbon backbone, *e.g.*, oxygen, nitrogen or sulfur atoms.

The terms "polycyclyl," "multicycle" or "polycyclic radical" refer to two or more cyclic rings (*e.g.*, cycloalkyls, cycloalkenyls, cycloalkynyls, aryls and/or heterocyclyls) in which two or more carbons are common to two adjoining rings, *e.g.*, the rings are  
20 "fused rings". Rings that are joined through non-adjacent atoms are termed "bridged" rings. Each of the rings of the polycycle can be substituted with such substituents as described above, as for example, halogen, hydroxyl, alkylcarbonyloxy, arylcarbonyloxy, alkoxycarbonyloxy, aryloxy carbonyloxy, carboxylate, alkylcarbonyl, alkoxycarbonyl, aminocarbonyl, alkylthiocarbonyl, alkoxyl, phosphate, phosphonato, phosphinato,  
25 cyano, amino (including alkyl amino, dialkylamino, arylamino, diarylamino, and alkylaryl amino), acylamino (including alkylcarbonylamino, arylcarbonylamino, carbamoyl and ureido), amidino, imino, sulfhydryl, alkylthio, arylthio, thiocarboxylate, sulfates, sulfonato, sulfamoyl, sulfonamido, nitro, trifluoromethyl, cyano, azido, heterocyclyl, alkyl, alkylaryl, or an aromatic or heteroaromatic moiety. Examples of  
30 "multicyclic" moieties include steroids, such as, for example, cholesterol.

The term "heteroatom" as used herein means an atom of any element other than carbon or hydrogen. Preferred heteroatoms are nitrogen, oxygen, sulfur and phosphorus.

Suitable alkanoyl groups include groups having 1 to about 4 or 5 carbonyl groups. Suitable aroyl groups include groups having one or more carbonyl groups as a substituent to an aryl group such as phenyl or other carbocyclic aryl. Suitable alkaroyl groups have one or more alkylcarbonyl groups as a substituent to an aryl group such as phenylacetyl and the like. Suitable carbocyclic aryl groups have 6 or more carbons such as phenyl, naphthyl and the like. Suitable aryloyl groups are carbocyclic aryl groups that are substituted with one or more carbonyl groups, typically 1 or 2 carbonyl groups.

Prodrugs are compounds which are converted *in vivo* to active forms (see, *e.g.*, R.B. Silverman, 1992, "The Organic Chemistry of Drug Design and Drug Action", Academic Press, Chp. 8). Prodrugs can be used to alter the biodistribution (*e.g.*, to allow compounds which would not typically enter the reactive site of the protease) or the pharmacokinetics for a particular compound. For example, a hydroxyl group, can be esterified, *e.g.*, with a carboxylic acid group to yield an ester. When the ester is administered to a subject, the ester is cleaved, enzymatically or non-enzymatically, reductively or hydrolytically, to reveal the hydroxyl group.

The language "prodrug moiety" includes moieties which can be metabolized *in vivo* to yield an active compound. For example, the term includes moieties which can modify certain functional groups of the substituted tetracycline compounds, such as, but not limited to, hydroxyl groups and amino groups. In an embodiment, the prodrugs moieties are metabolized *in vivo* by esterases or by other mechanisms to hydroxyl groups, amino, amido or other groups which allow the substituted tetracycline compound to perform its intended function. Examples of prodrugs and their uses are well known in the art (See, *e.g.*, Berge et al. (1977) "Pharmaceutical Salts", *J. Pharm. Sci.* 66:1-19). Some prodrugs can be prepared *in situ* during the final isolation and purification of the compounds, or by separately reacting the purified compound in its free acid form or hydroxyl with a suitable esterifying agent. Hydroxyl groups, for example, can be converted into esters *via* treatment with a carboxylic acid (see, for example, Scheme 3). Examples of prodrug moieties include substituted and unsubstituted, branch or unbranched lower alkyl ester moieties, (*e.g.*, propionoic acid esters), lower alkenyl esters, di-lower alkyl-amino lower-alkyl esters (*e.g.*, dimethylaminoethyl ester), acylamino lower alkyl esters (*e.g.*, acetyloxymethyl ester), acyloxy lower alkyl esters (*e.g.*, pivaloyloxymethyl ester), aryl esters (phenyl ester),

- 30 -

aryl-lower alkyl esters (*e.g.*, benzyl ester), substituted (*e.g.*, with methyl, halo, or methoxy substituents) aryl and aryl-lower alkyl esters, amides, lower-alkyl amides, di-lower alkyl amides, and hydroxy amides. Preferred prodrug moieties for hydroxyl groups are propionoic acid esters and acyl esters. Amino or amido groups can be  
5 modified by methods known in the art to form Schiff bases and other prodrugs which may or may not be metabolized *in vivo*.

It will be noted that the structure of some of the compounds of this invention includes asymmetric carbon atoms. It is to be understood accordingly that the isomers arising from such asymmetry (*e.g.*, all enantiomers and diastereomers) are included  
10 within the scope of this invention, unless indicated otherwise. Such isomers can be obtained in substantially pure form by classical separation techniques and by stereochemically controlled synthesis.

The invention also features a method for treating a tetracycline compound responsive state in a subject, by administering to the subject a substituted tetracycline  
15 compound of the invention. Preferably, an effective amount of the substituted tetracycline compound is administered. In an embodiment, the substituted tetracycline compound is of formula (I). The invention includes methods of treating a tetracycline compound responsive state using any one of the compounds described above or found in Table 2, below.

20 The term "subject" includes any animal or plant which is capable of being treated or may obtain some benefit from the administration of a substituted tetracycline compound of the invention. The term also include animals (*e.g.*, birds, reptiles, fish, mammals, (*e.g.*, cows, pigs, sheep, horses, cows, dogs, cats, squirrels, bears, monkeys, chimpanzees, gorillas, goats, ferrets, and, preferably, humans). The subject may be  
25 currently suffering from the tetracycline compound responsive state or may be at risk of suffering from the tetracycline compound responsive state. In an embodiment, the subject may be immunocompromised, *e.g.*, suffering from AIDS, undergoing or recovering from chemotherapy, or have an immune disorder.

The language "tetracycline compound responsive state" includes state which can  
30 be treated, prevented, or otherwise ameliorated by the administration of a substituted tetracycline compound of the invention. Tetracycline compound responsive states include bacterial infections (including those which are resistant to other tetracycline

- 31 -

compounds), cancer, diabetes, and other states for which tetracycline compounds have been found to be active (see, for example, U.S. Patent Nos. 5,789,395; 5,834,450; and 5,532,227). Compounds of the invention can be used to prevent or control important mammalian and veterinary diseases such as diarrhea, urinary tract infections, infections of skin and skin structure, ear, nose and throat infections, wound infection, mastitis and the like. In addition, methods for treating neoplasms using tetracycline compounds of the invention are also included (van der Bozert et al., *Cancer Res.*, 48:6686-6690 (1988)).

Bacterial infections may be caused by a wide variety of gram positive and gram negative bacteria. The compounds of the invention are useful as antibiotics against organisms which are resistant to other tetracycline compounds. The antibiotic activity of the tetracycline compounds of the invention may be determined using the method discussed in Example 2, or by using the in vitro standard broth dilution method described in Waitz, J.A., *National Commission for Clinical Laboratory Standards, Document M7-A2*, vol. 10, no. 8, pp. 13-20, 2<sup>nd</sup> edition, Villanova, PA (1990).

The tetracycline compounds may also be used to treat infections traditionally treated with tetracycline compounds such as, for example, rickettsiae; a number of gram-positive and gram-negative bacteria; and the agents responsible for lymphogranuloma venereum, inclusion conjunctivitis, psittacosis. The substituted tetracycline compounds may be used to treat infections of *pneumococci*, *Salmonella*, *E. coli*, *S. aureus*, *E. hirae* or *E. faecalis*. In one embodiment, the substituted tetracycline compound is used to treat a bacterial infection that is resistant to other unsubstituted tetracycline antibiotic compounds (e.g., tetracycline compounds such as doxycycline, minocycline, sancycline, or tetracycline). In another embodiment, the substituted tetracycline compounds of the invention are less cytotoxic to the subject as compared to unsubstituted tetracycline compounds, such that the substituted tetracycline compounds may be given at a higher dosage without being fatal or excessively toxic to the subject. The substituted tetracycline compound of the invention may be administered with a pharmaceutically acceptable carrier.

Examples of compounds of the invention which may advantageously be used in the methods of the invention include substituted tetracycline compounds of formula (I),



- 32 -

- as well as compounds described in Table 2. Examples of compounds of the invention include: Doxycycline 9-carbamic acid 9*H*-fluoren-9-ylmethyl ester;
- (9-(Naphthyn-1-yl) doxycycline urea;
- 9-(3-Methyl-1-butyl) doxycycline urea;
- 5 9-Phenyl doxycycline urea;
- 9-*t*-Butyl doxycycline urea;
- Fmoc 9-amino doxycycline;
- 9-(4'-Chloro-2'-trifluoromethylphenyl) doxycycline urea;
- 9-(4'-Fluorophenyl) doxycycline carbamate;
- 10 9-(4'-Methoxyphenyl) doxycycline carbamate;
- 9-BOC amino doxycycline;
- 9-(Phenylthiazolyl) amino doxycycline;
- 9-(Ethylthiazolyl) amino doxycycline;
- (4-Fluorophenylthiazolyl) amino doxycycline;
- 15 9-(4'-Methoxyphenylthiazolyl) amino doxycycline;
- 9-(3'-Nitrophenylthiazolyl) amino doxycycline;
- 9-(4'-Methyl, 5'-phenylthiazolyl) amino doxycycline;
- 9-Neopentyl minocycline carbamate;
- 9-(Phenylthiazolyl) amino sancycline;
- 20 9-(Adamantylthiazolyl) amino doxycycline;
- 9-(Naphthyn-1-yl urea) Doxycycline 5-propanoic acid ester;
- Doxycycline 9-Thiocarbamic acid 9*H*-fluoren-9-ylmethyl ester;
- (9-(Naphthyn-1-yl) doxycycline thiourea;
- 9-(3-methyl-1-butyl) doxycycline thiourea;
- 25 9-Phenyl doxycycline thiourea;
- 9-*t*-Butyl doxycycline thiourea;
- 9-(4'-Chloro-2'-trifluoromethylphenyl) doxycycline thiourea;
- 9-(4'-Fluorophenyl) doxycycline thiocarbamate;
- 9-(4-Methoxyphenyl) doxycycline thiocarbamate;
- 30 9-Neopentyl minocycline thiocarbamate;
- 9-(Naphthyn-1-yl) doxycycline thiourea 5-propanoic acid ester;
- Minocycline 9-carbamic acid 9*H*-fluoren-9-ylmethyl ester;

- 33 -

- (9-(Naphthyn-1-yl) minocycline urea;
- 9-(3-Methyl-1-butyl) minocycline urea;
- 9-Phenyl doxycycline urea;
- 9-t-Butyl minocycline urea;
- 5 Fmoc 9-amino minocycline;
- 9-(4'-Chloro-2'-trifluoromethylphenyl) minocycline urea;
- 9-(4'-Fluorophenyl) minocycline carbamate;
- 9-(4'-Methoxyphenyl) minocycline carbamate;
- 9-BOC amino minocycline;
- 10 9-(Phenylthiazolyl) amino minocycline;
- 9-(Ethylthiazolyl) amino minocycline;
- (4'-Fluorophenylthiazolyl) amino minocycline;
- 9-(4'-Methoxyphenylthiazolyl) amino minocycline;
- 9-9-(3'-Nitrophenylthiazolyl) amino minocycline;
- 15 9-(4'-Methyl, 5'-phenylthiazolyl) amino doxycycline;
- 9-Neopentyl doxycycline carbamate;
- 9-(Phenylthiazolyl) amino minocycline;
- 9-(Adamantylthiazolyl) amino minocycline;
- Minocycline 9-thiocarbamic acid 9*H*-fluoren-9-ylmethyl ester;
- 20 (9-(Naphthyn-1-yl) minocycline thiourea;
- 9-(3'-Methyl-1-butyl) minocycline thiourea;
- 9-Phenyl minocycline thiourea;
- 9-t-Butyl minocycline thiourea;
- 9-(4'-Fluorophenyl) minocycline thiocarbamate;
- 25 9-(4'-Methoxyphenyl) minocycline thiocarbamate;
- 9-Neopentyl doxycycline thiocarbamate;
- 9-(2'-Bromoethyl) doxycycline carbamate;
- 9-(n-Pentyl) minocycline carbamate;
- 9-(4'-Benzoylbenzoyl) amino doxycycline;
- 30 7-(3'-Nitrophenylthiazolyl) amino sancycline;
- 9-(3'-Ethoxycarbonylthiazolyl) amino doxycycline;
- 7-(4'-Methylphenyl) sancycline carbamate;

- 34 -

- 9-(4'-Trifluoromethoxyphenyl) minocycline urea;
- 9-(3', 5'-diperfluorophenyl) minocycline thiourea;
- 9-Prop-2'-enyl minocycline carbamate;
- 9-(4'-Chloro, 2'-nitrophenyl) minocycline urea;
- 5 9-Ethyl minocycline carbamate;
- 9-n-Butyl minocycline carbamate;
- 9-n-But-3-enyl minocycline carbamate;
- Doxycycline 7-carbamic acid 7*H*-fluoren-7-ylmethyl ester;
- 7-(Naphthyn-1-yl) doxycycline urea;
- 10 7-(3-Methyl-1-butyl) doxycycline urea;
- 7-Phenyl doxycycline urea;
- 7-t-Butyl doxycycline urea;
- 7-Fmoc amino doxycycline;
- 7-(4'-Chloro-2-trifluoromethylphenyl) doxycycline urea;
- 15 7-(4'-Fluorophenyl) doxycycline carbamate;
- 7-(4'-Methoxyphenyl) doxycycline carbamate;
- 7-BOC amino doxycycline;
- 7-(3'-Phenylthiazolyl) amino doxycycline;
- 7-(3'-Ethylthiazolyl) amino doxycycline;
- 20 7-(4''-Fluorophenylthiazolyl) amino doxycycline;
- 7-(4''-Methoxyphenylthiazolyl) amino doxycycline;
- 7-(Phenylthiazolylamino)-sancycline;
- 7-(3'-Nitrophenylthiazolyl) amino doxycycline;
- 7-(4'-Methyl, 5'-phenylthiazolyl) amino doxycycline;
- 25 7-(Adamantylthiazolyl) amino doxycycline;
- Doxycycline 7-thiocarbamic acid 7*H*-fluoren-7-ylmethyl ester;
- 7-(Naphthyn-1-yl) doxycycline thiourea;
- 7-(3-Methyl-1-butyl) doxycycline thiourea;
- 7-Phenyl amino doxycycline thiourea;
- 30 7-t-butyl amino doxycycline thiourea;
- 7-(4'-Chloro-2'-trifluoromethylphenyl) doxycycline thiourea;
- 7-(4'-Fluorophenyl) doxycycline thiocarbamate;

- 35 -

7-(4'-Methoxyphenyl) doxycycline thiocarbamate;  
7-(Naphthyn-1-yl) doxycycline urea 5-propanoic acid ester;  
7-(Naphthyn-1-yl) doxycycline thiourea 5-propanoic acid ester;  
9-i-Butyl minocycline carbamate, and pharmaceutically acceptable salts and prodrugs  
5 thereof.

The language "effective amount" of the substituted tetracycline compound is that amount necessary or sufficient to treat or prevent a tetracycline compound responsive state. The effective amount can vary depending on such factors as the size and weight of the subject, the type of illness, or the particular substituted tetracycline compound.  
10 For example, the choice of the substituted tetracycline compound can affect what constitutes an "effective amount". One of ordinary skill in the art would be able to study the aforementioned factors and make the determination regarding the effective amount of the substituted tetracycline compound without undue experimentation.

The invention also pertains to methods of treatment against microorganism  
15 infections and associated diseases. The methods include administration of an effective amount of one or more substituted tetracycline compounds to a subject. The subject can be either a plant or, advantageously, an animal, *e.g.*, a mammal, *e.g.*, a human.

In the therapeutic methods of the invention, one or more substituted tetracycline compounds of the invention may be administered alone to a subject, or more typically a  
20 compound of the invention will be administered as part of a pharmaceutical composition in mixture with conventional excipient, *i.e.*, pharmaceutically acceptable organic or inorganic carrier substances suitable for parenteral, oral or other desired administration and which do not deleteriously react with the active compounds and are not deleterious to the recipient thereof.

25 In one embodiment, the invention pertains to pharmaceutical compositions which comprise one or more substituted tetracycline compounds of the invention, as described above. The invention pertains to pharmaceutical compositions which comprise any of the substituted tetracycline compounds described in this application. For example, the invention pertains to pharmaceutical compositions which comprise  
30 substituted tetracycline compounds of both Formula (I) and Table 2. Other examples of substituted tetracycline compounds of the invention which may be included in the pharmaceutical compositions of the invention include, but are not limited to:

- 36 -

- Doxycycline 9-carbamic acid 9*H*-fluoren-9-ylmethyl ester;  
(9-(Naphthyn-1-yl) doxycycline urea;  
9-(3-Methyl-1-butyl) doxycycline urea;  
9-Phenyl doxycycline urea;  
5 9-*t*-Butyl doxycycline urea;  
FMOC 9-amino doxycycline;  
9-(4'-Chloro-2'-trifluoromethylphenyl) doxycycline urea;  
9-(4'-Fluorophenyl) doxycycline carbamate;  
9-(4'-Methoxyphenyl) doxycycline carbamate;  
10 9-BOC amino doxycycline;  
9-(Phenylthiazolyl) amino doxycycline;  
9-(Ethylthiazolyl) amino doxycycline;  
(4-Fluorophenylthiazolyl) amino doxycycline;  
9-(4'-Methoxyphenylthiazolyl) amino doxycycline;  
15 9-(3'-Nitrophenylthiazolyl) amino doxycycline;  
9-(4'-Methyl, 5'-phenylthiazolyl) amino doxycycline;  
9-Neopentyl minocycline carbamate;  
9-(Phenylthiazolyl) amino sancycline;  
9-(Adamantylthiazolyl) amino doxycycline;  
20 9-(Naphthyn-1-yl urea) Doxycycline 5-propanoic acid ester;  
Doxycycline 9-Thiocarbamic acid 9*H*-fluoren-9-ylmethyl ester;  
(9-(Naphthyn-1-yl) doxycycline thiourea;  
9-(3-methyl-1-butyl) doxycycline thiourea;  
9-Phenyl doxycycline thiourea;  
25 9-*t*-Butyl doxycycline thiourea;  
9-(4'-Chloro-2'-trifluoromethylphenyl) doxycycline thiourea;  
9-(4'-Fluorophenyl) doxycycline thiocarbamate;  
9-(4-Methoxyphenyl) doxycycline thiocarbamate;  
9-Neopentyl minocycline thiocarbamate;  
30 9-(Naphthyn-1-yl) doxycycline thiourea 5-propanoic acid ester;  
Minocycline 9-carbamic acid 9*H*-fluoren-9-ylmethyl ester;  
(9-(Naphthyn-1-yl) minocycline urea;

- 37 -

- 9-(3-Methyl-1-butyl) minocycline urea;
- 9-Phenyl doxycycline urea;
- 9-t-Butyl minocycline urea;
- Fmoc 9-amino minocycline;
- 5 9-(4'-Chloro-2'-trifluoromethylphenyl) minocycline urea;
- 9-(4'-Fluorophenyl) minocycline carbamate;
- 9-(4'-Methoxyphenyl) minocycline carbamate;
- 9-BOC amino minocycline;
- 9-(Phenylthiazolyl) amino minocycline;
- 10 9-(Ethylthiazolyl) amino minocycline;
- (4'-Fluorophenylthiazolyl) amino minocycline;
- 9-(4'-Methoxyphenylthiazolyl) amino minocycline;
- 9-9-(3'-Nitrophenylthiazolyl) amino minocycline;
- 9-(4'-Methyl, 5'-phenylthiazolyl) amino doxycycline;
- 15 9-Neopentyl doxycycline carbamate;
- 9-(Phenylthiazolyl) amino minocycline;
- 9-(Adamantylthiazolyl) amino minocycline;
- Minocycline 9-thiocarbamic acid 9*H*-fluoren-9-ylmethyl ester;
- (9-(Naphthyl-1-yl) minocycline thiourea;
- 20 9-(3'-Methyl-1-butyl) minocycline thiourea;
- 9-Phenyl minocycline thiourea;
- 9-t-Butyl minocycline thiourea;
- 9-(4'-Fluorophenyl) minocycline thiocarbamate;
- 9-(4'-Methoxyphenyl) minocycline thiocarbamate;
- 25 9-Neopentyl doxycycline thiocarbamate;
- 9-(2'-Bromoethyl) doxycycline carbamate;
- 9-(n-Pentyl) minocycline carbamate;
- 9-(4'-Benzoylbenzoyl) amino doxycycline;
- 7-(3'-Nitrophenylthiazolyl) amino sancycline;
- 30 9-(3'-Ethoxycarbonylthiazolyl) amino doxycycline;
- 7-(4'-Methylphenyl) sancycline carbamate;
- 9-(4'-Trifluoromethoxyphenyl) minocycline urea;

- 38 -

- 9-(3', 5'-diperfluorophenyl) minocycline thiourea;  
9-Prop-2'-enyl minocycline carbamate;  
9-(4'-Chloro, 2'-nitrophenyl) minocycline urea;  
9-Ethyl minocycline carbamate;
- 5 9-n-Butyl minocycline carbamate;  
9-n-But-3-enyl minocycline carbamate;  
Doxycycline 7-carbamic acid 7*H*-fluoren-7-ylmethyl ester;  
7-(Naphthyn-1-yl) doxycycline urea;  
7-(3-Methyl-1-butyl) doxycycline urea;
- 10 7-Phenyl doxycycline urea;  
7-t-Butyl doxycycline urea;  
7-Fmoc amino doxycycline;  
7-(4'-Chloro-2-trifluoromethylphenyl) doxycycline urea;  
7-(4'-Fluorophenyl) doxycycline carbamate;
- 15 7-(4'-Methoxyphenyl) doxycycline carbamate;  
7-BOC amino doxycycline;  
7-(3'-Phenylthiazolyl) amino doxycycline;  
7-(3'-Ethylthiazolyl) amino doxycycline;  
7-(4''-Fluorophenylthiazolyl) amino doxycycline;
- 20 7-(4''-Methoxyphenylthiazolyl) amino doxycycline;  
7-(Phenylthiazolylamino)-sancycline;  
7-(3'-Nitrophenylthiazolyl) amino doxycycline;  
7-(4'-Methyl, 5'-phenylthiazolyl) amino doxycycline;  
7-(Adamantylthiazolyl) amino doxycycline;
- 25 Doxycycline 7-thiocarbamic acid 7*H*-fluoren-7-ylmethyl ester;  
7-(Naphthyn-1-yl) doxycycline thiourea;  
7-(3-Methyl-1-butyl) doxycycline thiourea;  
7-Phenyl amino doxycycline thiourea;  
7-t-butyl amino doxycycline thiourea;
- 30 7-(4'-Chloro-2'-trifluoromethylphenyl) doxycycline thiourea;  
7-(4'-Fluorophenyl) doxycycline thiocarbamate;  
7-(4'-Methoxyphenyl) doxycycline thiocarbamate;

- 39 -

7-(Naphthyn-1-yl) doxycycline urea 5-propanoic acid ester;  
7-(Naphthyn-1-yl) doxycycline thiourea 5-propanoic acid ester;  
9-i-Butyl minocycline carbamate, and pharmaceutically acceptable salts and prodrugs thereof.

5           The language "pharmaceutically acceptable carrier" includes substances capable of being coadministered with the substituted tetracycline compound(s), and which allow the substituted tetracycline compound to perform its intended function, *e.g.*, treat or prevent a tetracycline compound responsive state. Suitable pharmaceutically acceptable carriers include but are not limited to water, salt solutions, alcohol, vegetable oils,  
10 polyethylene glycols, gelatin, lactose, amylose, magnesium stearate, talc, silicic acid, viscous paraffin, perfume oil, fatty acid monoglycerides and diglycerides, petroethral fatty acid esters, hydroxymethyl-cellulose, polyvinylpyrrolidone, *etc.* The pharmaceutical preparations can be sterilized and if desired mixed with auxiliary agents, *e.g.*, lubricants, preservatives, stabilizers, wetting agents, emulsifiers, salts for  
15 influencing osmotic pressure, buffers, colorings, flavorings and/or aromatic substances and the like which do not deleteriously react with the active compounds of the invention.

          The substituted tetracycline compounds of the invention that are basic in nature are capable of forming a wide variety of salts with various inorganic and organic acids. The acids that may be used to prepare pharmaceutically acceptable acid addition salts of  
20 the tetracycline compounds of the invention that are basic in nature are those that form non-toxic acid addition salts, *i.e.*, salts containing pharmaceutically acceptable anions, such as the hydrochloride, hydrobromide, hydroiodide, nitrate, sulfate, bisulfate, phosphate, acid phosphate, isonicotinate, acetate, lactate, salicylate, citrate, acid citrate, tartrate, pantothenate, bitartrate, ascorbate, succinate, maleate, gentisinate, fumarate,  
25 gluconate, glucaronate, saccharate, formate, benzoate, glutamate, methanesulfonate, ethanesulfonate, benzenesulfonate, p-toluenesulfonate and palmoate [*i.e.*, 1,1'-methylene-bis-(2-hydroxy-3-naphthoate)] salts. Although such salts must be pharmaceutically acceptable for administration to a subject, *e.g.*, a mammal, it is often desirable in practice to initially isolate a tetracycline compound of the invention from  
30 the reaction mixture as a pharmaceutically unacceptable salt and then simply convert the latter back to the free base compound by treatment with an alkaline reagent and subsequently convert the latter free base to a pharmaceutically acceptable acid addition



- 40 -

salt. The acid addition salts of the base compounds of this invention are readily prepared by treating the base compound with a substantially equivalent amount of the chosen mineral or organic acid in an aqueous solvent medium or in a suitable organic solvent, such as methanol or ethanol. Upon careful evaporation of the solvent, the  
5 desired solid salt is readily obtained.

The substituted tetracycline compounds of the invention that are acidic in nature are capable of forming a wide variety of base salts. The chemical bases that may be used as reagents to prepare pharmaceutically acceptable base salts of those substituted tetracycline compounds of the invention that are acidic in nature are those that form  
10 non-toxic base salts with such compounds. Such non-toxic base salts include, but are not limited to those derived from such pharmaceutically acceptable cations such as alkali metal cations (*e.g.*, potassium and sodium) and alkaline earth metal cations (*e.g.*, calcium and magnesium), ammonium or water-soluble amine addition salts such as N-methylglucamine-(meglumine), and the lower alkanolammonium and other base salts of  
15 pharmaceutically acceptable organic amines. The pharmaceutically acceptable base addition salts of tetracycline compounds of the invention that are acidic in nature may be formed with pharmaceutically acceptable cations by conventional methods. Thus, these salts may be readily prepared by treating the tetracycline compound of the invention with an aqueous solution of the desired pharmaceutically acceptable cation and  
20 evaporating the resulting solution to dryness, preferably under reduced pressure. Alternatively, a lower alkyl alcohol solution of the substituted tetracycline compound of the invention may be mixed with an alkoxide of the desired metal and the solution subsequently evaporated to dryness.

The preparation of other tetracycline compounds of the invention not specifically  
25 described in the foregoing experimental section can be accomplished using combinations of the reactions described above that will be apparent to those skilled in the art.

The substituted tetracycline compounds of the invention and pharmaceutically acceptable salts thereof can be administered via either the oral, parenteral or topical  
30 routes. In general, these compounds are most desirably administered in effective dosages, depending upon the weight and condition of the subject being treated and the particular route of administration chosen. Variations may occur depending upon the

- 41 -

species of the subject being treated and its individual response to said medicament, as well as on the type of pharmaceutical formulation chosen and the time period and interval at which such administration is carried out.

The pharmaceutical compositions of the invention may be administered alone or  
5 in combination with other known compositions for treating tetracycline responsive states in a mammal. Preferred mammals include pets (*e.g.*, cats, dogs, ferrets, *etc.*), farm animals (cows, sheep, pigs, horses, goats, *etc.*), lab animals (rats, mice, monkeys, *etc.*), and primates (chimpanzees, humans, gorillas). The language "in combination with" a known composition is intended to include simultaneous administration of the  
10 composition of the invention and the known composition, administration of the composition of the invention first, followed by the known composition and administration of the known composition first, followed by the composition of the invention. Any of the therapeutically composition known in the art for treating tetracycline responsive states can be used in the methods of the invention.

15 The substituted tetracycline compounds of the invention may be administered alone or in combination with pharmaceutically acceptable carriers or diluents by any of the routes previously mentioned, and the administration may be carried out in single or multiple doses. For example, the novel therapeutic agents of this invention can be administered advantageously in a wide variety of different dosage forms, *i.e.*, they may  
20 be combined with various pharmaceutically acceptable inert carriers in the form of tablets, capsules, lozenges, troches, hard candies, powders, sprays, creams, salves, suppositories, jellies, gels, pastes, lotions, ointments, aqueous suspensions, injectable solutions, elixirs, syrups, and the like. Such carriers include solid diluents or fillers, sterile aqueous media and various non-toxic organic solvents, *etc.* Moreover, oral  
25 pharmaceutical compositions can be suitably sweetened and/or flavored. In general, the therapeutically-effective compounds of this invention are present in such dosage forms at concentration levels ranging from about 5.0% to about 70% by weight.

For oral administration, tablets containing various excipients such as microcrystalline cellulose, sodium citrate, calcium carbonate, dicalcium phosphate and  
30 glycine may be employed along with various disintegrants such as starch (and preferably corn, potato or tapioca starch), alginic acid and certain complex silicates, together with granulation binders like polyvinylpyrrolidone, sucrose, gelatin and acacia. Additionally,

- 42 -

lubricating agents such as magnesium stearate, sodium lauryl sulfate and talc are often very useful for tableting purposes. Solid compositions of a similar type may also be employed as fillers in gelatin capsules; preferred materials in this connection also include lactose or milk sugar as well as high molecular weight polyethylene glycols.

5 When aqueous suspensions and/or elixirs are desired for oral administration, the active ingredient may be combined with various sweetening or flavoring agents, coloring matter or dyes, and, if so desired, emulsifying and/or suspending agents as well, together with such diluents as water, ethanol, propylene glycol, glycerin and various like combinations thereof.

10 For parenteral administration (including intraperitoneal, subcutaneous, intravenous, intradermal or intramuscular injection), solutions of a substituted tetracycline compound of the present invention in either sesame or peanut oil or in aqueous propylene glycol may be employed. The aqueous solutions should be suitably buffered (preferably pH greater than 8) if necessary and the liquid diluent first rendered  
15 isotonic. These aqueous solutions are suitable for intravenous injection purposes. The oily solutions are suitable for intraarticular, intramuscular and subcutaneous injection purposes. The preparation of all these solutions under sterile conditions is readily accomplished by standard pharmaceutical techniques well known to those skilled in the art. For parenteral application, examples of suitable preparations include solutions,  
20 preferably oily or aqueous solutions as well as suspensions, emulsions, or implants, including suppositories. Substituted tetracycline compounds may be formulated in sterile form in multiple or single dose formats such as being dispersed in a fluid carrier such as sterile physiological saline or 5% saline dextrose solutions commonly used with injectables.

25 Additionally, it is also possible to administer the substituted tetracycline compounds of the present invention topically when treating inflammatory conditions of the skin. Examples of methods of topical administration include transdermal, buccal or sublingual application. For topical applications, therapeutic compounds can be suitably admixed in a pharmacologically inert topical carrier such as a gel, an ointment, a lotion  
30 or a cream. Such topical carriers include water, glycerol, alcohol, propylene glycol, fatty alcohols, triglycerides, fatty acid esters, or mineral oils. Other possible topical carriers are liquid petrolatum, isopropylpalmitate, polyethylene glycol, ethanol 95%,

- 43 -

polyoxyethylene monolaurate 5% in water, sodium lauryl sulfate 5% in water, and the like. In addition, materials such as anti-oxidants, humectants, viscosity stabilizers and the like also may be added if desired.

For enteral application, particularly suitable are tablets, dragees or capsules  
5 having talc and/or carbohydrate carrier binder or the like, the carrier preferably being lactose and/or corn starch and/or potato starch. A syrup, elixir or the like can be used wherein a sweetened vehicle is employed. Sustained release compositions can be formulated including those wherein the active component is protected with differentially degradable coatings, *e.g.*, by microencapsulation, multiple coatings, *etc.*

10 In addition to treatment of human subjects, the therapeutic methods of the invention also will have significant veterinary applications, *e.g.* for treatment of livestock such as cattle, sheep, goats, cows, swine and the like; poultry such as chickens, ducks, geese, turkeys and the like; horses; and pets such as dogs and cats. Also, the compounds of the invention may be used to treat non-animal subjects, such as plants.

15 It will be appreciated that the actual preferred amounts of active compounds used in a given therapy will vary according to the specific compound being utilized, the particular compositions formulated, the mode of application, the particular site of administration, *etc.* Optimal administration rates for a given protocol of administration can be readily ascertained by those skilled in the art using conventional dosage  
20 determination tests conducted with regard to the foregoing guidelines.

In general, compounds of the invention for treatment can be administered to a subject in dosages used in prior tetracycline therapies. See, for example, the *Physicians' Desk Reference*. For example, a suitable effective dose of one or more compounds of the invention will be in the range of from 0.01 to 100 milligrams per kilogram of body  
25 weight of recipient per day, preferably in the range of from 0.1 to 50 milligrams per kilogram body weight of recipient per day, more preferably in the range of 1 to 20 milligrams per kilogram body weight of recipient per day. The desired dose is suitably administered once daily, or several sub-doses, *e.g.* 2 to 5 sub-doses, are administered at appropriate intervals through the day, or other appropriate schedule.

30 It will also be understood that normal, conventionally known precautions will be taken regarding the administration of tetracyclines generally to ensure their efficacy under normal use circumstances. Especially when employed for therapeutic treatment

- 44 -

of humans and animals *in vivo*, the practitioner should take all sensible precautions to avoid conventionally known contradictions and toxic effects. Thus, the conventionally recognized adverse reactions of gastrointestinal distress and inflammations, the renal toxicity, hypersensitivity reactions, changes in blood, and impairment of absorption  
5 through aluminum, calcium, and magnesium ions should be duly considered in the conventional manner.

### **Exemplification of the Invention**

The following example illustrates various methods of synthesizing 9- and 7-  
10 substituted tetracycline compounds in accordance with the invention. Other compounds of the invention can be synthesized using methods described herein and/or methods known in the art.

#### **Example 1: Synthesis of 9- Amino-Substituted Tetracycline Compounds**

15 To an ice cold solution of doxycycline (2 g, 4.15 mmol) in 30 ml of concentrated H<sub>2</sub>SO<sub>4</sub>, potassium nitrate (0.5g; 1.2 eq) was added portion-wise. The reaction mixture was stirred for 1 1/2 hours. The acid solution was then added to ~200 ml of ice water. The precipitated yellow material was filtered. The filtered material was dissolved in methanol. After the methanol was evaporated, the product was extracted with n-butanol.  
20 The organic phase was washed with saturated NaHCO<sub>3</sub> twice and the solvent removed *in vacuo*.

The resulting product was dissolved in 50 ml of methanol and 1 ml of concentrated HCl and hydrogenated on Pd/C to yield the 9- and 7- aminodoxycycline positional isomers as an off-yellow solid. The isomers can be purified by HPLC and  
25 other techniques known in the art.

#### **9-Aminodoxycycline**

(9-amino-6- $\alpha$ -deoxy-5-hydroxytetracycline)

MS(M + H):460 <sup>1</sup>HNMR (CD<sub>3</sub>OD):  $\delta$  7.62 (d, 1H, H-8); 7.14 (d, 1H, H-7); 4.42 (s, 1H,  
30 H-4); 3.6 (dd, 1H, H-5); 2.98, 2.90 (each s, each 3H, NMe<sub>2</sub>); 2.84(d, 1H, H-4a); 2.72 (m, 1H, H-6); 2.59 (dd, 1H, H-5a); 1.51 (d, 3H, CH<sub>3</sub>-C<sub>6</sub>).

- 45 -

**General Synthesis of mixed urea of 9- or 7- Amino-6- $\alpha$ -tetracycline compound:**

In one portion, 1.2 equivalents of an isocyanate was added to a solution of 9-amino tetracycline compound in DMF and two equivalents of diisopropylethylamine. The reaction mixture was stirred at room temperature for several hours (usually 4 hours  
5 to overnight). The desired product was isolated by C18 reverse-phase column chromatography. The 7- amino tetracycline compound urea can be synthesized using the similar methodology with the 7-amino tetracycline compound as the starting material.

The following compounds were made using the above procedure.

10

Compound B: 9-Aminonaphthyl Doxycycline urea

(1-Naphthyl, 9-amino-6- $\alpha$ -deoxy-5-hydroxy-tetracycline mixed urea)

<sup>1</sup>HNMR (CD<sub>3</sub>OD):  $\delta$  7.9 (d, 1H, H-8); 7.8-7.4 (m, 7H, aryl); 6.9 (bd, 1H, H-7); 4.42 (s, 1H, H-4); 3.6 (d, 1H, H-5); 2.88, 2.77 (each s, each 3H, NMe<sub>2</sub>); 2.84 (d, 1H, H-4a); 2.72  
15 (m, 1H, H-6); 2.59 (dd, 1H, H-5a); 1.31 (d, 3H, CH<sub>3</sub>-C<sub>6</sub>). MS(M + H): calc. 629.63; Found 629.16.

Compound E: 9-Aminophenyl Doxycycline Urea

(Phenyl, 9-amino- $\alpha$ -deoxy-5-hydroxy-tetracycline mixed urea)

<sup>1</sup>HNMR (CD<sub>3</sub>OD):  $\delta$  8.25 (d, 1H, H-8); 7.45 (d, 2H, aryl); 7.29 (m, 2H, aryl); 7.0 (d, 1H, aryl); 6.9 (d, 1H, H-7); 4.25 (s, 1H, H-4); 3.6 (dd, 1H, H-5); 2.85 (bs, 6H, NMe<sub>2</sub>);  
20 2.84 (d, 1H, H-4a); 2.72 (m, 1H, H-6); 2.58 (dd, 1H, H-5a); 1.54 (d, 3H, CH<sub>3</sub>-C<sub>6</sub>). MS(M + H): calc. 579.57; Found 579.15.

Compound F: 9-Amino-t-butyl doxycycline urea

(Tert-butyl, 9-amino-6- $\alpha$ -deoxy-5-hydroxy-tetracycline mixed urea)

<sup>1</sup>HNMR (CD<sub>3</sub>OD):  $\delta$  8.1 (d, 1H, H-8); 6.84 (d, 1H, H-7); 4.31 (s, 1H, H-4); 3.55 (dd, 1H, H-5); 2.91 (bs, 6H, NMe<sub>2</sub>); 2.71-2.57 (m, 3H, H-4a, H-6, H-5a); 1.51 (d, 3H, CH<sub>3</sub>-C<sub>6</sub>); 1.36 (s, 9H, tert butyl); MS(M + H): calc. 559.58; Found: 559.19.

30

- 46 -

Compound I: 9-(4'-Chloro, 2'-trifluoromethylphenyl amino)-Doxycycline urea

4-Chloro, 2-trifluoromethylphenyl, 9-amino-6- $\alpha$ -deoxy-5-hydroxy-tetracycline mixed urea

<sup>1</sup>HNMR (CD<sub>3</sub>OD):  $\delta$  8.28 (d, 1H, H-8); 7.89 (d, 1H, aryl); 7.66 (s, 1H, aryl); 7.58 (d, 1H, aryl); 6.93 (d, 1H, H-7); 4.42 (s, 1H, H-4); 3.56 (dd, 1H, H-5); 2.98 & 2.90 (each s, 3H, NMe<sub>2</sub>); 2.84 (d, 2H, H-4a); 2.72 (m, 1H, H-6); 2.56 (dd, 1H, H-5a); 1.52 (d, 3H, CH<sub>3</sub>-C<sub>6</sub>). MS(M + H): calc.

Compound AJ: 9-(3,5-Bis (trifluoromethylphenyl) amino) Doxycycline urea

(3,5-bis (trifluoromethyl phenyl), 9-amino-6- $\alpha$ -deoxy-5-hydroxy-tetracycline mixed urea)

**General Synthesis of Carbamates of 9- or 7- amino-doxycycline:**

In a single portion, 1.2 equivalents of chloroformate was added to a solution of 9-aminodoxycycline in DMF in the presence of two equivalent of diisopropylethylamine. The reaction mixture was stirred at room temperature for several hours. The desired product was isolated by C18 reverse-phase column chromatography. The 7- amino tetracycline compound carbamate can be synthesized using the similar methodology with the 7-amino tetracycline compound as the starting material. The following carbamates were synthesized using the general synthesis outlined above.

Compound A: FMOC-9-Amino Doxycycline

(N-Fluorenylmethyloxycarbonyl 9-amino-6- $\alpha$ -deoxy-5-hydroxy-tetracycline)

<sup>1</sup>HNMR (CD<sub>3</sub>OD)  $\delta$  7.9 (bd, 1H, H-8), 7.69 (d, 2H, aryl); 7.56 (m, 2H, aryl); 7.29 (m, 4H, aryl); 6.8 (d, 1H, H-7); 4.35 (d, 2H, CH<sub>2</sub>) 4.30 (s, 1H, H-4); 4.15 (m, 1H, CH); 3.5 (dd, 1H, H-5); 2.85 (bd, 6H, NMe<sub>2</sub>); 2.83 (d, 1H, H-4a); 2.73 (m, 1H, H-6a); 2.57 (dd, 1H, H-5a); 1.40 (d, 3H, CH<sub>3</sub>-C<sub>6</sub>). MS(M + H): calc. 682.69; Found: 682.

- 47 -

**Compound K: 9-(Fluorophenyl) Doxycycline Carbamate**(N-p-Fluorophenyloxycarbonyl 9-amino-6- $\alpha$ -deoxy-5-hydroxy tetracycline)

<sup>1</sup>HNMR (CD<sub>3</sub>OD)  $\delta$ : 7.92 (d, 1H, H-8); 7.11-6.98 (m, 4H, aryl); 6.85 (d, 1H, H-7); 4.34 (s, 1H, H-4); 3.42 (dd, 1H, H-5); 2.86 (bd, 6H, NMe<sub>2</sub>); 2.83 (d, 1H, H-5a); 2.72 (m, 1H, H-6); 2.56 (dd, 1H, H-5a); 1.43 (d, 3H, CH<sub>3</sub>-C<sub>6</sub>). MS(M + H): calc. 598.55; Found: 598.50

**Compound L: 9-(4-Methoxyphenyl) Doxycycline Carbamate**(N-p-methoxyphenyloxycarbonyl 9-amino-6- $\alpha$ -deoxy-5-hydroxy tetracycline)

<sup>1</sup>HNMR (CD<sub>3</sub>OD):  $\delta$  7.92 (bd, 1H, H-8); 6.97 (d, 2H, aryl); 6.82 (m, 3H, H-7 and aryl); 4.36 (s, 1H, H-4); 3.66 (s, 3H, OMe); 3.4 (d, 1H, H-5); 2.86 (bd, 6H, NMe<sub>2</sub>); 2.83 (d, 1H, H-4a); 2.78 (m, 1H, H-6); 2.56 (dd, 1H, H-5a); 1.43 (d, 3H, CH<sub>3</sub>-C<sub>6</sub>). MS(M + H): calc. 610.58; Found: 610.50.

**Compound M: 9-BOC-Amino Doxycycline**(N-tert-butyloxycarbonyl 9-amino-6- $\alpha$ -deoxy-5-hydroxy-tetracycline)

<sup>1</sup>HNMR (CD<sub>3</sub>OD):  $\delta$  8.04 (d, 1H, H-8); 6.92 (d, 1H, H-7); 4.05 (s, 1H, H-4); 3.62 (dd, 1H, H-5); 2.82 (bs, 6H, NMe<sub>2</sub>); 2.83 (d, 1H, H-4a); 2.74 (m, 1H, H-6); 2.57 (dd, 1H, H-5a); 1.52 (bs, 12H, CH<sub>3</sub>-C<sub>6</sub> + tert-butyl); MS(M + H): calc. 560.57; found: 560.16.

**Compound AP: 9-Neopentyl Minocycline Carbamate**(N-neopentyloxycarbonyl 9-amino-6- $\alpha$ -deoxy-5-hydroxy-tetracycline)

<sup>1</sup>HNMR (CD<sub>3</sub>OD):  $\delta$  7.9 (d, 1H, H-8); 6.9 (d, 1H, H-7); 4.36 (s, 1H, H-4); 3.77 (s, 2H, neopentyl CH<sub>2</sub>); 3.6 (dd, 1H, H-5); 2.88, 2.81 (bs, 6H, NMe<sub>2</sub>); 2.84 (d, 1H, H-4a); 2.72 (m, 1H, H-6); 2.59 (dd, 1H, H-5a); 1.45 (d, 3H, CH<sub>3</sub>-C<sub>6</sub>); 0.89 (s, 9H, neopentyl CH<sub>3</sub>). MS(M + H): calc. 587.63; Found: 587.5.

**Synthesis of 2-Aminothiazole Derivatives of Tetracycline Compounds**

Fluorenylmethyloxycarbonyl chloride (1.80 g; 5 mmol) was dissolved in 10 ml of ethyl acetate. This solution was added drop-wise to a suspension of potassium thiocyanate (1.2 eq) in 10 ml of ethyl acetate at 0° and under nitrogen atmosphere. The



- 48 -

reaction mixture was left stirring overnight. The reaction mixture was then filtered over celite pad to remove residual salts, and the ethyl acetate was removed *in vacuo*. The crude yellow material was used to synthesize the compounds below.

5    Compound AT: 9-FMOC-amino Doxycycline thiocarboxamide

(3-(Fluorenylmethyloxycarbonyl)-1-(9-amino-6- $\alpha$ -deoxy-5-hydroxy tetracycline)-thio carboxamide)

To 300 mg (0.65 mmol) of 9-amino doxycycline in 3 ml of DMF and in the presence of 227  $\mu$ l (2 eq) of diisopropylethylamine, was added in one portion of 182 mg  
10 of fluorenylmethoxy-carbonyl isothiocyanate in 1 ml of DMF. The reaction mixture was stirred at room temperature for 5 hours. The desired product was isolated through C18 reverse-phase column chromatography.

$^1\text{H}$ NMR ( $\text{CD}_3\text{OD}$ ):  $\delta$  8.82 (d, 1H, H-8); 7.82 (d, 2H, aryl); 7.72 (d, 2H, aryl); 7.4 (m, 4H, aryl); 6.92 (d, 1H, H-7); 4.56 (d, 2H,  $\text{CH}_2$ ); 4.44 (s, 1, H-4); 4.30 (m, 1H,  
15 CH); 3.6 (dd, 1H, H-5); 2.98 (bd, 6H,  $\text{NMe}_2$ ); 2.84 (d, 1H, H-4a); 2.73 (m, 1H, H-6); 2.56 (dd, 1H, H-5a); 1.54 (d, 3H,  $\text{CH}_3\text{-C}_6$ ). MS(M + H): calc. 741.78; Found: 741.28.

6- $\alpha$ -deoxy-5-hydroxy-tetracycline thio-urea

300 mg (0.405 mmol) of 3- (fluorenylmethyloxycarbonyl)-1- (9-amino-6- $\alpha$ -  
20 deoxy-5-hydroxy tetracycline)- thio carboxamide was deblocked in a solution of 2% piperidine, 2% DBU in DMF at room temperature. The solvent was then evaporated *in vacuo* after acidification with concentrated HCl. The residue was dissolved in 1 ml of MeOH and added dropwise to 100 ml of cold ethyl acetate. The precipitated yellow solid was filtered and dried.  $^1\text{H}$ NMR ( $\text{CD}_3\text{OD}$ ):  $\delta$  7.90 (d, 1H, H-8); 6.95 (d, 1H, H-7);  
25 4.48 (s, 1H, H-4); 3.57 (dd, 1H, H-5); 3.04, 2.92 (two s, each 3H,  $\text{NMe}_2$ ); 2.84 (d, 1H, H-4a); 2.7 (m, 1H, H-6); 2.6 (dd, 1H, H-5a); 1.54 (d, 3H,  $\text{CH}_3\text{-C}_6$ ). MS(M + H): calc. 519.54; Found: 519.20.

**General Synthesis of 7- or 9- (2' -thiazolyl amino) Tetracycline Compounds**

30    In one portion, the appropriate  $\alpha$ -bromo ketone was added to a tetracycline compound thiourea in a mixture of DMF/Dioxane (3:1) and an equivalent amount of diisopropylethylamine. The reaction mixture was left stirring overnight. The thiazole

product was isolated through C18 reverse-phase column chromatography. The following thiazole compounds were synthesized using the method described above.

Compound N: 9-(4'-Phenyl thiazolyl)-amino Doxycycline

5 2 [9 (amino-6- $\alpha$ -deoxy-5-hydroxy tetracycline)]-4- phenyl thiazole:

$^1\text{HNMR}$  ( $\text{CD}_3\text{OD}$ ):  $\delta$  8.25 (d, 1H, H-8); 7.8 (d, 2H, aryl); 7.45 (m, 3H, aryl); 7.1 (s, 1H, vinyl); 7.09 (d, 1H, H-7); 4.46 (s, 1H, H-4); 3.6 (dd, 1H, H-5); 2.91 & 2.88 (Two s, each 3H,  $\text{NMe}_2$ ); 2.84 (d, 1H, H-4a); 2.7 (m, 1H, H-6); 2.57 (dd, 1H, H-5a); 1.6 (d, 3H,  $\text{CH}_3\text{-C}_6$ ). MS(M + H): calc. 619.66; Found: 619.19.

10

Compound O: 9-(4'-Ethyl thiazolyl)-amino Doxycycline

2 [9-(amino-6- $\alpha$ -deoxy-5-hydroxy tetracycline)-4-ethyl] thiazole

15  $^1\text{HNMR}$  ( $\text{CD}_3\text{OD}$ ):  $\delta$  7.9 (d, 1H, H-8); 7.05 (d, 1H, H-7); 6.55 (s, 1H, vinyl); 4.46 (s, 1H, H-4); 3.57 (dd, 1H, H-5a); 2.96 (bs, 6H,  $\text{NMe}_2$ ); 2.87 (d, 1H, H-4a); 2.7 (m, 1H, H-6); 2.6 (m, 3H, H-5a and  $\text{CH}_2$  of the ethyl); 1.59 (d, 3H,  $\text{CH}_3\text{-C}_6$ ); 1.28 (d, 3H,  $\text{CH}_3$  of the ethyl); MS(M + H): calc. 571.62; Found: 571.2.

Compound Q: 9-(4-Methoxyphenylthiazolyl)-amino Doxycycline

(2 [(9-amino-6- $\alpha$ -deoxy-5-hydroxy tetracycline)]-4- (4-methoxyphenyl) thiazole)

20  $^1\text{HNMR}$  ( $\text{CD}_3\text{OD}$ ):  $\delta$  7.94 (d, 1H, H-8); 7.68 (d, 2H, aryl); 7.10 (d, 1H, H-7); 7.06 (d, 2H, aryl); 4.49 (s, 1H, H-4); 3.86 (s, 3H, OMe); 3.56 (dd, 1H, H-5); 3.0 & 2.94 (two s, each 3H,  $\text{NMe}_2$ ); 2.87 (d, 1H, H-4a); 2.73 (m, 1H, H-6); 1.63 (d, 3H,  $\text{CH}_3\text{-C}_6$ ). MS(M + H): calc. 649.68; Found: 649.15.

25 Compound R: 9-(3-Nitrophenylthiazolyl)-amino Doxycycline

(2 [9-(amino-6- $\alpha$ -deoxy-5-hydroxy tetracycline)]-4-(3-nitrophenyl) thiazole)

30  $^1\text{HNMR}$  ( $\text{CD}_3\text{OD}$ ):  $\delta$  8.6 (m, 2H, aryl); 8.2 (d, 1H, H-8); 8.1 (d, 1H, aryl); 7.6 (m, 1H, aryl); 7.3 (s, 1H, vinyl); 6.9 (d, 1H, H-7); 4.44 (s, 1H, H-4); 3.57 (dd, 1H, H-5); 3.0 & 2.91 (two s, each 3H,  $\text{NMe}_2$ ); 2.84 (s, 1H, H-4a); 2.7 (m, 1H, H-6); 2.57 (dd, 1H, H-5a); 1.56 (d, 3H,  $\text{CH}_3\text{-C}_6$ ). MS(M + H): calc. 664.66; Found: 664.60.

- 50 -

Compound S: 9-(4-Methyl-5-phenylthiazolyl)-amino Doxycycline(2 [9(amino-6- $\alpha$ -deoxy-5-hydroxy tetracycline)]-4-phenyl-5-methyl thiazole)

<sup>1</sup>HNMR (CD<sub>3</sub>OD):  $\delta$  7.98 (d, 1H, H-8); 7.6 - 7.4 (m, 5H, aryl); 7.05 (d, 1H, H-7); 4.46 (s, 1H, H-4); 3.57 (dd, 1H, H-5); 2.95 (bs, 6H, NMe<sub>2</sub>); 2.87 (d, 1H, H-4a); 2.7 (m, 1H, H-6); 2.6 (dd, 1H, H-5a); 2.36 (s, 3H, CH<sub>3</sub>); 1.57 (d, 3H, CH<sub>3</sub>-C<sub>6</sub>). MS(M + H): calc. 633.68; Found: 633. 61.

Compound U: (9-(N,N-Dimethyl Glycyl)-Doxycycline)

NN-Dimethylglycine (1.2 mmol) is dissolved in DMF (5 mL) and *O*-Benzotriazol-1-yl-*N, N, N', N'*-tetramethyluronium hexafluorophosphate (HBTU, 1.2 mmol) is added. The solution is then stirred for 5 minutes at room temperature. To this solution, 9-amino doxycycline (1 mmol) is added, followed by the addition of diisopropylethyl amine (DIEA, 1.2 mmol). The reaction is then stirred at room temperature for 2 hours. The solvent, DMF, is removed under vacuum. The crude material is dissolved in 5 mL of MeOH and filtered using autovials and purified using preparative HPLC. The structure of the product is characterized using <sup>1</sup>H NMR, HPLC, and MS.

**Example 2: *In vitro* Minimum Inhibitory Concentration (MIC) Assay**

The following assay was used to determine the efficacy of tetracycline compounds against common bacteria (*E. coli*, *S. aureus*, *E. hirae*, and *E. faecalis*). 2 mg of each compound was dissolved in 100  $\mu$ L of DMSO. The solution was then added to cation-adjusted Mueller Hinton broth (CAMHB), which resulted in a final compound concentration of 200  $\mu$ g per mL. The tetracycline compound solutions were diluted to 50  $\mu$ L volumes, with a test compound concentration of .098  $\mu$ g/mL. Optical density (OD) determinations were made from fresh log-phase broth cultures of the test strains. Dilutions were made to achieve a final cell density of about  $5 \times 10^5$  CFU/mL.

50  $\mu$ L of the cell suspensions were added to each well of the microtiter plates. The final cell density was approximately  $5 \times 10^5$  CFU/mL. These plates were incubated at 35°C in an ambient air incubator for approximately 18 hr.

The plates were read with a microplate reader and were visually inspected when necessary. The MIC is defined as the lowest concentration of the tetracycline compound that inhibits growth. In Table 2, \* indicates good inhibition of the growth of a particular

- 51 -







organism, \*\* indicates inhibition of growth at a lower concentration, and \*\*\* indicates very good inhibition of growth. Certain substituted tetracycline compounds of the invention had MIC's less than about 10 µg/ml. Other substituted tetracycline compounds of the invention had MIC's of less than about 5 µg/mL, and still other  
5 compounds had MIC's less than about 1 µg/mL.







### EQUIVALENTS







Those skilled in the art will recognize, or be able to ascertain using no more than routine experimentation, numerous equivalents to the specific procedures described  
10 herein. Such equivalents are considered to be within the scope of the present invention and are covered by the following claims. The contents of all references, patents, and patent applications cited throughout this application are hereby incorporated by reference. The appropriate components, processes, and methods of those patents, applications and other documents may be selected for the present invention and  
15 embodiments thereof.

TABLE 2







ID	STRUCTURE	NAME	S. aureus	E. hirae	E. coli	E. faecalis
A		Fmoc-9-amino Doxycycline	***	***	*	***
B		9-Naphth-1-yl doxycycline urea	**	**	**	***
C		9-(3'-methyl butyl) Doxycycline urea	**	NT	**	**
D		9-(Naphth-1-yl) doxycycline urea 5-propanoic acid ester	**	NT	**	**
E		9-Phenyl Doxycycline urea	**	*	**	***
F		9-t-butyl doxycycline urea	**	**	*	**







I		9-(4'-chloro, 2'-perfluoromethyl)-doxycycline urea	**	NT	**	**
K		9-(4'-Fluorophenyl) doxycycline carbamate	*	*	*	*
L		9-(4'-Methoxyphenyl) doxycycline carbamate	*	*	*	*
M		9-BOC-amino doxycycline	**	**	**	**
N		9-(phenylthiazolyl)amino doxycycline	***	NT	*	***
O		9-(ethyl thiazolyl) amino doxycycline	***	NT	*	**







P		9-(4'-Fluorophenylthiazolyl)-amino doxycycline	*	NT	*	**
Q		9-(4'-Methoxyphenylthiazolyl)-amino doxycycline	**	NT	*	**
R		9-(3'-Nitrophenylthiazolyl)-amino doxycycline	***	***	*	***
S		9-(4'-Methyl, 3'-phenylthiazolyl)-amino doxycycline	**	NT	*	***
T		9-(Adamantylthiazolyl)-amino doxycycline	*	*	*	*
U		9-(N,N-Dimethylglycyl)-doxycycline	*	*	*	NT







V		9-(N,N-Dimethylamino-glycyl) minocycline	NT	NT	NT	NT
W		9-propyl doxycycline urea	**	*	**	**
X		9-(4'-chlorophenyl sulfonyl) doxycycline urea	**	*	*	**
Y		9-valine doxycycline urea	*	*	*	*
Z		9-Cholesterol doxycycline carbamate	*	*	*	*
AA		9-(2,2'-dimethylpropyl) doxycycline carbamate	**	**	**	**















AB		9-Tolyl doxycycline carbamate	*	*	*	*
AC		9-Naphthyl doxycycline thiourea	**	**	*	**
AD		9-(4'-Chloromethylphenyl) doxycycline urea	**	**	*	**
AE		9-(4'-Chloromethyl, 2'-trifluoromethylphenyl) doxycycline urea	**	**	**	**
AF		9-(4-chloro,2-trifluoromethylphenyl) minocycline urea	**	***	*	***
AG		9-Naphthyl minocycline urea	**	**	**	**








AH		9-Dansyl-phenyl doxycycline thiourea	*	*	*	*
AI		9-DABF-phenyl doxycycline thiourea	**	***	*	***
AJ		9(3,5-Bis (trifluoromethyl)phenyl doxycycline urea	*	*	*	*
AK		8-Chloro-8-N,N- Dimethylamino glycyI doxycycline	*	*	*	*
AL		9-Formylamino doxycycline	**	*	**	***
AM		9-Propenyl doxycycline carbamate	*	*	**	***





AN		9-Bromoethyl doxycycline carbamate	**	**	**	***
AO		9-Acetamide doxycycline	*	*	*	**
AP		9-(2,2-dimethyl propyl) minocycline carbamate	**	**	*	**
AQ		9-Isopropenyl minocycline carbamate	*	*	*	*
AR		9-(4'- Benzoylbenzoyl)amino doxycycline	**	**	*	**
AS		9-BOC-amino minocycline	**	**	*	**

AT		3-FMOC-9-amino doxycycline-thiourea	NT	NT	NT	NT
AU		9-(Phenylthiazolyl)amino doxycycline	***	**	*	**
AV		9-(4-Diethylaminophenyl thiazolyl)-amino doxycycline	*	NT	*	*
AW		9-(Biphenylthiazolyl) amino-doxycycline	*	*	*	*
AX		7-(3'- Nitrophenylthiazolyl) amino sancycline	**	***	*	***
AY		9- (Ethoxycarbonylthiazolyl) amino doxycycline	**	**	*	*

AZ		7-(4'-Methoxyphenyl) sancycline carbamate	**	**	*	**
BA		7-(4'-Methylphenyl) Sancycline Carbamate	***	***	*	**
BB		9-(Naphth-1-yl) minocycline thiourea	**	**	*	**
BC		9-(Phenyl) minocycline urea	**	**	**	**
BD		9-(4'-phenyl) minocycline urea	**	**	*	**
BE		9-(4'-methoxyphenyl) minocycline urea	*	**	**	**

BF		9-(4'-trifluoromethoxyphenyl) minocycline urea	**	**	**	**
BG		9-(Benzoyl) Minocycline urea	**	**	*	**
BH		9-(2', 5'-diperfluoromethyl phenyl) minocycline thiourea	**	**	**	**
BI		9-(4'-Nitrophenyl)-minocycline thiourea	**	**	*	**
BJ		9-(prop-2-enyl) minocycline carbamate	**	**	*	**
BK		9-(4'-Fluoro-3'-nitrophenyl)-minocycline urea	**	**	*	**


BL		9-(4'-Chloro-2'-nitrophenyl)-minocycline urea	**	**	**	**
BM		9-(3'-Fluorophenyl)-minocycline urea	**	**	*	**
BN		9-Phenyl minocycline thiocarbamate	*	*	*	*
BO		9-(4'-Bromophenyl)-minocycline carbamate	*	*	*	*
BP		9-(4'-Chlorophenyl)-minocycline carbamate	*	*	*	*
BQ		9-(Ethyl)-minocycline carbamate	***	**	**	**
BR		9-(n-Butyl)-minocycline carbamate	**	**	**	**

BS		9-(Ethyl)-minocycline thiocarbamate	**	**	*	*
BT		9-(But-3-enyl) minocycline carbamate	**	**	**	**
BU		9-(Phenyl)-minocycline carbamate	*	*	*	*
BV		9-(Isobutyl)-minocycline carbamate	**	**	**	**



## CLAIMS

1. A substituted tetracycline compound, wherein said compound is of the formula:



5

(I)

wherein:

X is  $\text{CHC}(\text{R}^{13}\text{Y}'\text{Y})$ ,  $\text{CR}^6\text{R}^6$ , S,  $\text{NR}^6$ , or O;

$\text{R}^2$  is hydrogen, alkyl, alkynyl, alkoxy, alkylthio, alkylsulfinyl, alkylsulfonyl, alkylamino, arylalkyl, aryl, heterocyclic, heteroaromatic or a prodrug moiety;

10

$\text{R}^4$  and  $\text{R}^{4'}$  are each hydrogen, alkyl, alkenyl, alkynyl, alkoxy, alkylthio, alkylsulfinyl, alkylsulfonyl, alkylamino, arylalkyl, aryl, heterocyclic, heteroaromatic or a prodrug moiety;

$\text{R}^{2'}$ ,  $\text{R}^3$ ,  $\text{R}^{10}$ ,  $\text{R}^{11}$  and  $\text{R}^{12}$  are each hydrogen or a pro-drug moiety;

15

$\text{R}^5$  is hydrogen, hydroxyl, or a prodrug moiety;

$\text{R}^6$ ,  $\text{R}^{6'}$ , and  $\text{R}^8$  are each independently hydrogen, alkyl, alkenyl, alkynyl, aryl, alkoxy, alkylthio, alkylsulfinyl, alkylsulfonyl, alkylamino, arylalkyl, or halogen;

$\text{R}^7$  is hydrogen, dialkylamino, heteroaryl-amino, or  $\text{NR}^{7c}\text{C}(=\text{W}')\text{WR}^{7a}$ ;

$\text{R}^{13}$  is hydrogen, hydroxy, alkyl; alkenyl; alkynyl; alkoxy; alkylthio;

20 alkylsulfinyl; alkylsulfonyl; alkylamino; or an arylalkyl;

$\text{Y}'$  and  $\text{Y}$  are each independently hydrogen; halogen; hydroxyl; cyano, sulfhydryl; amino; alkyl; alkenyl; alkynyl; alkoxy; alkylthio; alkylsulfinyl; alkylsulfonyl; alkylamino; or an arylalkyl;

$\text{R}^9$  is hydrogen,  $\text{NR}^{9c}\text{C}(=\text{Z}')\text{ZR}^{9a}$ , or heteroaryl-amino;

25 Z is  $\text{CR}^{9d}\text{R}^{9e}$ ,  $\text{NR}^{9b}$ , or O;

$\text{Z}'$  is O or S;

$\text{R}^{9a}$ ,  $\text{R}^{9b}$ ,  $\text{R}^{9c}$ ,  $\text{R}^{9d}$ , and  $\text{R}^{9e}$  are each independently hydrogen, alkyl, alkenyl, alkynyl, alkoxy, alkylthio, alkylsulfinyl, alkylsulfonyl, arylsulfonyl,

- 65 -

alkoxycarbonyl, arylcarbonyl, alkylamino, arylalkyl, aryl, heterocyclic, heteroaromatic, absent, or a prodrug moiety, and  $R^{9d}$  and  $R^{9e}$  may be linked to form a ring;

W is  $CR^{7d}R^{7e}$ ,  $NR^{7b}$  or O;

W' is O or S; and

- 5  $R^{7a}$ ,  $R^{7b}$ ,  $R^{7c}$ ,  $R^{7d}$ , and  $R^{7e}$  are each independently hydrogen, alkyl, alkenyl, alkynyl, alkoxy, alkylthio, alkylsulfinyl, arylsulfonyl, alkoxycarbonyl, arylcarbonyl, alkylamino, arylalkyl, aryl, heterocyclic, heteroaromatic, absent, or a prodrug moiety, and  $R^{7d}$  and  $R^{7e}$  may be linked to form a ring;
- and pharmaceutically acceptable salts thereof, provided that at least one
- 10 of  $R^9$  is not hydrogen when  $R^7$  is hydrogen or dialkylamino.


2. The compound of claim 1, wherein  $R^2$ ,  $R^{2'}$ ,  $R^3$ ,  $R^8$ ,  $R^{10}$ ,  $R^{11}$ , and  $R^{12}$  are each hydrogen.
- 15 3. The compound of claim 1 or 2, wherein  $R^4$  and  $R^{4'}$  are each alkyl.
4. The compound of claim 3, wherein  $R^4$  and  $R^{4'}$  are each methyl
5. The compound of claim 4, wherein said compound is a derivative of tetracycline,
- 20 minocycline, sancycline, doxycycline, chlortetracycline, oxytetracycline, demeclocycline, or methacycline.
6. The compound of any one of claims 1-5, wherein  $R^5$  is hydrogen.
- 25 7. The compound of claim 6, wherein X is  $CH_2$ , and  $R^7$  is hydrogen.
8. The compound of claim 6, wherein X is  $CH_2$ , and  $R^7$  is  $N(Me)_2$ .
9. The compound of anyone of claims 1-5, wherein  $R^5$  is hydroxyl or a prodrug
- 30 moiety, and X is  $CHR^6$ .
10. The compound of claim 9, wherein  $R^5$  is hydroxyl and  $R^6$  is  $CH_3$ .

- 66 -

11. The compound of any one of claims 1-10, wherein  $R^9$  is  $NR^{9c}C(=Z')ZR^{9a}$ .
12. The compound of claim 11, wherein  $R^{9c}$  is hydrogen.
- 5 13. The compound of claim 11 or 12, wherein  $Z'$  is oxygen.
14. The compound of any one of claims 11-13, wherein  $Z$  is  $NR^{9b}$ .
15. The compound of any one of claims 11-13, wherein  $Z$  is oxygen.
- 10 16. The compound of claim 11 or 12, wherein  $Z'$  is sulfur.
17. The compound of anyone of claims 11, 12, or 16, wherein  $Z$  is  $NR^{9b}$ .
- 15 18. The compound of anyone of claims 11, 12, or 16, wherein  $Z$  is oxygen.
19. The compound of anyone of claims 1-18, wherein  $R^{9a}$  is selected from the group consisting of alkyl, alkenyl, alkynyl, aryl, arylalkyl, heteroaromatic, and multicyclic.
- 20 20. The compound of claim 19, wherein  $R^{9a}$  is substituted or unsubstituted alkyl.
21. The compound of claim 20, wherein  $R^{9a}$  is substituted with one or more substituents selected from the group consisting of alkoxycarbonyl, amino, arylcarbonyl, halogen, hydroxy, alkylamino, alkoxy, or aryl.
- 25 22. The compound of claim 20, wherein  $R^{9a}$  is methyl, ethyl, t-butyl, n-butyl, i-butyl, or n-pentyl.
23. The compound of claim 21, wherein said alkyl is substituted with an aryl group.
- 30 24. The compound of claim 23, wherein said aryl group is phenyl.

- 67 -

25. The compound of claim 21, wherein said alkyl is substituted with one or more halogens.
26. The compound of claim 24, wherein said halogen is bromine.
- 5 27. The compound of claim 19, wherein  $R^{9a}$  is multicyclic.
28. The compound of claim 27, wherein  $R^{9a}$  is steroidyl.
- 10 29. The compound of claim 28, wherein  $R^{9a}$  is cholesterol.
30. The compound of claim 19, wherein  $R^{9a}$  is substituted or unsubstituted aryl.
31. The compound of claim 30, wherein said substituted or unsubstituted aryl is  
15 naphthyl.
32. The compound of claim 30, wherein said substituted or unsubstituted aryl is of the formula:



- 20 33. The compound of claim 30, wherein said substituted or unsubstituted aryl is phenyl.
34. The compound of claim 30 or 33, wherein said aryl is substituted with one or  
25 more substituents selected from the group consisting of alkyl, alkenyl, alkynyl, aryl, alkoxy, aryloxy, alkylcarbonyl, arylcarbonyl, alkoxycarbonyl, aryloxycarbonyl, amido, halogen, nitro, azo, alkyl sulfonyl, and arylsulfonyl.

- 68 -

35. The compound of claim 34, wherein said substituent is alkyl.
36. The compound of claim 35, wherein said alkyl is unsubstituted.
- 5 37. The compound of claim 35, wherein said alkyl is methyl.
38. The compound of claim 35, wherein said alkyl is substituted with one or more halogens.
- 10 39. The compound of claim 34, wherein said substituent is methoxy.
40. The compound of claim 34, wherein said substituent is selected from the group consisting of alkylcarbonyl, arylcarbonyl, alkoxycarbonyl, aryloxy, and amido.
- 15 41. The compound of any one of claims 1-10, wherein R<sup>9</sup> is heteroaryl-amino.
42. The compound of claim 41, wherein said heteroaryl is substituted or unsubstituted thioazolyl.
- 20 43. The compound of claim 42, wherein said heteroaryl is substituted thioazolyl.
44. The compound of claim 43, wherein said thiazolyl is substituted with a substituted or unsubstituted aryl.
- 25 45. The compound of claim 46, wherein said aryl is phenyl.
46. The compound of claim 44 or 45, wherein said aryl is substituted with one or more substituents selected from the group consisting of alkyl, alkenyl, alkynyl, aryl, alkoxy, aryloxy, alkylcarbonyl, arylcarbonyl, amido, trifluoromethyl, halogen, nitro, 30 azo, alkyl sulfonyl, and arylsulfonyl.
47. The compound of claim 46, wherein said substituent is nitro.

- 69 -

48. The compound of claim 46, wherein said substituent is alkyl.
49. The compound of claim 48, wherein said alkyl substituent is methyl.
50. The compound of claim 46, wherein said substituent is selected from the group consisting of alkylcarbonyl, arylcarbonyl, alkoxy carbonyl, aryloxy carbonyl, and amido.
51. The compound of claim 50, wherein said substituent is alkoxy carbonyl.
52. The compound of claim 51, wherein said substituent is ethoxy carbonyl.
53. The compound of claim 1, wherein said compound is selected from the group consisting of: Doxycycline 9-carbamic acid 9*H*-fluoren-9-ylmethyl ester;  
(9-(Naphthyn-1-yl) doxycycline urea;  
9-(3-Methyl-1-butyl) doxycycline urea;  
9-Phenyl doxycycline urea;  
9-*t*-Butyl doxycycline urea;  
Fmoc 9-amino doxycycline;  
9-(4'-Chloro-2'-trifluoromethylphenyl) doxycycline urea;  
9-(4'-Fluorophenyl) doxycycline carbamate;  
9-(4'-Methoxyphenyl) doxycycline carbamate;  
9-BOC amino doxycycline;  
9-(Phenylthiazolyl) amino doxycycline;  
9-(Ethylthiazolyl) amino doxycycline;  
9-(4-Fluorophenylthiazolyl) amino doxycycline;  
9-(4'-Methoxyphenylthiazolyl) amino doxycycline;  
9-(3'-Nitrophenylthiazolyl) amino doxycycline;  
9-(4'-Methyl, 5'-phenylthiazolyl) amino doxycycline;  
9-Neopentyl minocycline carbamate;  
9-(Phenylthiazolyl) amino sancycline;  
9-(Adamantylthiazolyl) amino doxycycline;  
9-(Naphthyn-1-yl urea) Doxycycline 5-propanoic acid ester;

- 70 -

- Doxycycline 9-Thiocarbamic acid 9*H*-fluoren-9-ylmethyl ester;  
(9-(Naphthyn-1-yl) doxycycline thiourea;  
9-(3-methyl-1-butyl) doxycycline thiourea;  
9-Phenyl doxycycline thiourea;  
5 9-*t*-Butyl doxycycline thiourea;  
9-(4'-Chloro-2'-trifluoromethylphenyl) doxycycline thiourea;  
9-(4'-Fluorophenyl) doxycycline thiocarbamate;  
9-(4-Methoxyphenyl) doxycycline thiocarbamate;  
9-Neopentyl minocycline thiocarbamate;  
10 9-(Naphthyn-1-yl) doxycycline thiourea 5-propanoic acid ester;  
Minocycline 9-carbamic acid 9*H*-fluoren-9-ylmethyl ester;  
(9-(Naphthyn-1-yl) minocycline urea;  
9-(3-Methyl-1-butyl) minocycline urea;  
9-Phenyl doxycycline urea;  
15 9-*t*-Butyl minocycline urea;  
Fmoc 9-amino minocycline;  
9-(4'-Chloro-2'-trifluoromethylphenyl) minocycline urea;  
9-(4'-Fluorophenyl) minocycline carbamate;  
9-(4'-Methoxyphenyl) minocycline carbamate;  
20 9-BOC amino minocycline;  
9-(Phenylthiazolyl) amino minocycline;  
9-(Ethylthiazolyl) amino minocycline;  
(4'-Fluorophenylthiazolyl) amino minocycline;  
9-(4'-Methoxyphenylthiazolyl) amino minocycline;  
25 9-9-(3'-Nitrophenylthiazolyl) amino minocycline;  
9-(4'-Methyl, 5'-phenylthiazolyl) amino doxycycline;  
9-Neopentyl doxycycline carbamate;



















- 71 -

54. The compound of claim 1, wherein said compound is selected from the group consisting of: 9-(Phenylthiazolyl) amino minocycline;  
9-(Adamantylthiazolyl) amino minocycline;  
Minocycline 9-thiocarbamic acid 9*H*-fluoren-9-ylmethyl ester;
- 5 (9-(Naphthyn-1-yl) minocycline thiourea;  
9-(3'-Methyl-1-butyl) minocycline thiourea;  
9-Phenyl minocycline thiourea;  
9-*t*-Butyl minocycline thiourea;  
9-(4'-Fluorophenyl) minocycline thiocarbamate;
- 10 9-(4'-Methoxyphenyl) minocycline thiocarbamate;  
9-Neopentyl doxycycline thiocarbamate;  
9-(2'-Bromoethyl) doxycycline carbamate;  
9-(*n*-Pentyl) minocycline carbamate;  
9-(4'-Benzoylbenzoyl) amino doxycycline;
- 15 7-(3'-Nitrophenylthiazolyl) amino sancycline;  
9-(3'-Ethoxycarbonylthiazolyl) amino doxycycline;  
7-(4'-Methylphenyl) sancycline carbamate;  
9-(4'-Trifluoromethoxyphenyl) minocycline urea;  
9-(3', 5'-diperfluorophenyl) minocycline thiourea;
- 20 9-Prop-2'-enyl minocycline carbamate;  
9-(4'-Chloro, 2'-nitrophenyl) minocycline urea;  
9-Ethyl minocycline carbamate;  
9-*n*-Butyl minocycline carbamate;  
9-*n*-But-3-enyl minocycline carbamate;
- 25 9-*i*-Butyl minocycline carbamate, and pharmaceutically acceptable salts and prodrugs thereof.

















- 72 -

55. The compound of claim 1, wherein said compound is selected from the group consisting of:



- 73 -



56. The compound of any one of claims 1-6 and 9-53, wherein  $R^7$  is  $NR^{7c}C(=W')WR^{7a}$ .

5 57. The compound of claim 56, wherein  $R^9$  is hydrogen.

- 74 -


58. The compound of claim 56 or 57, wherein  $R^{7c}$  is hydrogen.
59. The compound of any one of claims 56-58, wherein  $W'$  is oxygen.
- 5 60. The compound of any one of claims 56-58, wherein  $W'$  is sulfur
61. The compound of any one of claims 56-60, wherein  $W$  is  $NR^{7b}$ .
62. The compound of any one of claims 56-60, wherein  $W$  is oxygen.
- 10 63. The compound of any one of claims 56-62, wherein  $R^{7a}$  is selected from the group consisting of alkyl, alkenyl, alkynyl, aryl, arylalkyl, heteroaromatic, and multicyclic.
- 15 64. The compound of claim 63, wherein  $R^{7a}$  is substituted or unsubstituted alkyl.
65. The compound of claim 64, wherein said alkyl is substituted with an aryl group.
66. The compound of claim 63, wherein said substituted or unsubstituted aryl is  
20 phenyl.
67. The compound of claim 65 or 66, wherein said aryl is substituted with one or more substituents selected from the group consisting of alkyl, alkenyl, alkynyl, aryl, alkoxy, aryloxy, alkylcarbonyl, arylcarbonyl, alkoxycarbonyl, aryloxycarbonyl, amido,  
25 halogen, nitro, azo, alkyl sulfonyl, and arylsulfonyl.
68. The compound of claim 67, wherein said substituent is alkyl, alkoxy, or nitro.
69. The compound of any one of claims 1-6 and 9-52, wherein  $R^7$  is heteroaryl-  
30 amino.
70. The compound of claim 69, wherein  $R^9$  is hydrogen.

- 75 -

71. The compound of claim 69 or 70, wherein said heteroaryl is substituted or unsubstituted thioazolyl.
72. The compound of claim 71, wherein said thiazolyl is substituted with a  
5 substituted or unsubstituted aryl.
73. The compound of claim 72, wherein said aryl is phenyl.
74. The compound of claim 73, wherein said aryl is substituted with one or more  
10 substituents selected from the group consisting of alkyl, alkenyl, alkynyl, aryl, alkoxy, aryloxy, alkylcarbonyl, arylcarbonyl, amido, trifluoromethyl, halogen, nitro, azo, alkyl sulfonyl, and arylsulfonyl.
75. The compound of claim 74, wherein said substituent is nitro.  
15
76. The compound of claim 1, wherein said compound is selected from the group consisting of: Doxycycline 7-carbamic acid 7*H*-fluoren-7-ylmethyl ester;  
7-(Naphthyn-1-yl) doxycycline urea;  
7-(3-Methyl-1-butyl) doxycycline urea;  
20 7-Phenyl doxycycline urea;  
7-*t*-Butyl doxycycline urea;  
7-Fmoc amino doxycycline;  
7-(4'-Chloro-2-trifluoromethylphenyl) doxycycline urea;  
7-(4'-Fluorophenyl) doxycycline carbamate;  
25 7-(4'-Methoxyphenyl) doxycycline carbamate;  
7-BOC amino doxycycline;  
7-(3'Phenylthiazolyl) amino doxycycline;  
7-(3'-Ethylthiazolyl) amino doxycycline;  
7-(4"-Fluorophenylthiazolyl) amino doxycycline;  
30 7-(4"-Methoxyphenylthiazolyl) amino doxycycline;  
7-(Phenylthiazolylamino)-sancycline;  
7-(3'-Nitrophenylthiazolyl) amino doxycycline;

- 76 -

- 7-(4'-Methyl, 5'-phenylthiazolyl) amino doxycycline;  
 7-(Adamantylthiazolyl) amino doxycycline;  
 Doxycycline 7-thiocarbamic acid 7*H*-fluoren-7-ylmethyl ester;  
 7-(Naphthyn-1-yl) doxycycline thiourea;  
 5 7-(3-Methyl-1-butyl) doxycycline thiourea;  
 7-Phenyl amino doxycycline thiourea;  
 7-*t*-butyl amino doxycycline thiourea;  
 7-(4'-Chloro-2'-trifluoromethylphenyl) doxycycline thiourea;  
 7-(4'-Fluorophenyl) doxycycline thiocarbamate;  
 10 7-(4'-Methoxyphenyl) doxycycline thiocarbamate;  
 7-(Naphthyn-1-yl) doxycycline urea 5-propanoic acid ester;  
 7-(Naphthyn-1-yl) doxycycline thiourea 5-propanoic acid ester, and pharmaceutically acceptable salts thereof.
- 15 77. A method for treating a tetracycline responsive state in a mammal, comprising administering to said mammal a substituted tetracycline compound of formula (I):



(I)

wherein

- 20 X is  $\text{CHC}(\text{R}^{13}\text{Y}'\text{Y})$ ,  $\text{CR}^{6'}\text{R}^6$ , S,  $\text{NR}^6$ , or O;  
 $\text{R}^2$  is hydrogen, alkyl, alkynyl, alkoxy, alkylthio, alkylsulfinyl, alkylsulfonyl, alkylamino, arylalkyl, aryl, heterocyclic, heteroaromatic or a prodrug moiety;  
 $\text{R}^4$  and  $\text{R}^{4'}$  are each hydrogen, alkyl, alkenyl, alkynyl, alkoxy, alkylthio, alkylsulfinyl, alkylsulfonyl, alkylamino, arylalkyl, aryl, heterocyclic, heteroaromatic or  
 25 a prodrug moiety;  
 $\text{R}^{2'}$ ,  $\text{R}^3$ ,  $\text{R}^{10}$ ,  $\text{R}^{11}$  and  $\text{R}^{12}$  are each hydrogen or a pro-drug moiety;  
 $\text{R}^5$  is hydrogen, hydroxyl, or a prodrug moiety;

- 77 -

$R^6$ ,  $R^{6'}$ , and  $R^8$  are each independently hydrogen, alkyl, alkenyl, alkynyl, aryl, alkoxy, alkylthio, alkylsulfinyl, alkylsulfonyl, alkylamino, arylalkyl, or halogen;

$R^7$  is hydrogen, dialkylamino, heteroaryl-amino, or  $NR^{7c}C(=W')WR^{7a}$ ;

$R^{13}$  is hydrogen, hydroxy, alkyl; alkenyl; alkynyl; alkoxy; alkylthio; alkylsulfinyl; alkylsulfonyl; alkylamino; or an arylalkyl;

$Y'$  and  $Y$  are each independently hydrogen; halogen; hydroxyl; cyano, sulfhydryl; amino; alkyl; alkenyl; alkynyl; alkoxy; alkylthio; alkylsulfinyl; alkylsulfonyl; alkylamino; or an arylalkyl;

$R^9$  is hydrogen,  $NR^{9c}C(=Z')ZR^{9a}$ , or heteroaryl-amino;

$Z$  is  $CR^{9d}R^{9e}$ ,  $NR^{9b}$ , or O;

$Z'$  is O or S;

$R^{9a}$ ,  $R^{9b}$ ,  $R^{9c}$ ,  $R^{9d}$ , and  $R^{9e}$  are each independently hydrogen, alkyl, alkenyl, alkynyl, alkoxy, alkylthio, alkylsulfinyl, alkylsulfonyl, arylsulfonyl, alkoxycarbonyl, arylcarbonyl, alkylamino, arylalkyl, aryl, heterocyclic, heteroaromatic, absent, or a prodrug moiety, and  $R^{9d}$  and  $R^{9e}$  may be linked to form a ring;

$W$  is  $CR^{7d}R^{7e}$ ,  $NR^{7b}$  or O;

$W'$  is O or S; and

$R^{7a}$ ,  $R^{7b}$ ,  $R^{7c}$ ,  $R^{7d}$ , and  $R^{7e}$  are each independently hydrogen, alkyl, alkenyl, alkynyl, alkoxy, alkylthio, alkylsulfinyl, arylsulfonyl, alkoxycarbonyl, arylcarbonyl, alkylamino, arylalkyl, aryl, heterocyclic, heteroaromatic, absent, or a prodrug moiety, and  $R^{7d}$  and  $R^{7e}$  may be linked to form a ring;

and pharmaceutically acceptable salts thereof, provided that when  $R^9$  is not hydrogen when  $R^7$  is hydrogen or dialkylamino.

78. The method of claim 77, wherein said tetracycline compound is a compound of any one of claims 1-76.

79. The method of claim 77 or 78, wherein said tetracycline responsive state is a bacterial infection.


80. The method of claim 79, wherein said bacterial infection is associated with *E. coli*, *S. aureus*, *E. faecalis*, or *E. hirae*.

- 78 -

81. The method of claim 79 or 80, wherein said bacterial infection is resistant to unsubstituted tetracycline compounds.

82. The method of claim 77 or 78, wherein said tetracycline compound is  
5 administered with a pharmaceutically acceptable carrier.

83. A pharmaceutical composition comprising a therapeutically effective amount of a substituted tetracycline compound and a pharmaceutically acceptable carrier, wherein said substituted tetracycline is of the formula:



(I)

wherein:

X is  $\text{CHC}(\text{R}^{13}\text{Y}'\text{Y})$ ,  $\text{CR}^{6'}\text{R}^6$ , S,  $\text{NR}^6$ , or O;

$\text{R}^2$  is hydrogen, alkyl, alkynyl, alkoxy, alkylthio, alkylsulfinyl,  
15 alkylsulfonyl, alkylamino, arylalkyl, aryl, heterocyclic, heteroaromatic or a prodrug moiety;

$\text{R}^4$  and  $\text{R}^{4'}$  are each hydrogen, alkyl, alkenyl, alkynyl, alkoxy, alkylthio, alkylsulfinyl, alkylsulfonyl, alkylamino, arylalkyl, aryl, heterocyclic, heteroaromatic or a prodrug moiety;

20  $\text{R}^{2'}$ ,  $\text{R}^3$ ,  $\text{R}^{10}$ ,  $\text{R}^{11}$  and  $\text{R}^{12}$  are each hydrogen or a pro-drug moiety;

$\text{R}^5$  is hydrogen, hydroxyl, or a prodrug moiety;

$\text{R}^6$ ,  $\text{R}^{6'}$ , and  $\text{R}^8$  are each independently hydrogen, alkyl, alkenyl, alkynyl, aryl, alkoxy, alkylthio, alkylsulfinyl, alkylsulfonyl, alkylamino, arylalkyl, or halogen;

$\text{R}^7$  is hydrogen, dialkylamino, heteroaryl-amino, or  $\text{NR}^{7c}\text{C}(=\text{W}')\text{WR}^{7a}$ ;

25  $\text{R}^{13}$  is hydrogen, hydroxy, alkyl, alkenyl, alkynyl, alkoxy, alkylthio, alkylsulfinyl, alkylsulfonyl, alkylamino; or an arylalkyl;

$\text{Y}'$  and  $\text{Y}$  are each independently hydrogen; halogen; hydroxyl; cyano, sulfhydryl; amino; alkyl; alkenyl; alkynyl; alkoxy; alkylthio; alkylsulfinyl; alkylsulfonyl; alkylamino; or an arylalkyl;

- 79 -

$R^9$  is hydrogen,  $NR^{9c}C(=Z')ZR^{9a}$ , or heteroaryl-amino;

Z is  $CR^{9d}R^{9e}$ ,  $NR^{9b}$ , or O;

Z' is O or S;

- 5  $R^{9a}$ ,  $R^{9b}$ ,  $R^{9c}$ ,  $R^{9d}$ , and  $R^{9e}$  are each independently hydrogen, alkyl, alkenyl, alkynyl, alkoxy, alkylthio, alkylsulfinyl, alkylsulfonyl, arylsulfonyl, alkoxycarbonyl, arylcarbonyl, alkylamino, arylalkyl, aryl, heterocyclic, heteroaromatic, absent, or a prodrug moiety, and  $R^{9d}$  and  $R^{9e}$  may be linked to form a ring;

W is  $CR^{7d}R^{7e}$ ,  $NR^{7b}$  or O;

W' is O or S; and

- 10  $R^{7a}$ ,  $R^{7b}$ ,  $R^{7c}$ ,  $R^{7d}$ , and  $R^{7e}$  are each independently hydrogen, alkyl, alkenyl, alkynyl, alkoxy, alkylthio, alkylsulfinyl, arylsulfonyl, alkoxycarbonyl, arylcarbonyl, alkylamino, arylalkyl, aryl, heterocyclic, heteroaromatic, absent, or a prodrug moiety, and  $R^{7d}$  and  $R^{7e}$  may be linked to form a ring;
- and pharmaceutically acceptable salts thereof, provided that  $R^9$  is not
- 15 hydrogen, when  $R^7$  is dialkylamino or hydrogen.

84. The pharmaceutical composition of claim 83, wherein said substituted tetracycline compound is a compound of any of claims 1-76.

- 20 85. The pharmaceutical composition of claim 83 or 84, wherein said therapeutically effective amount is effective for treatment or prevention of a bacterial infection.

86. A method for synthesizing a 7- or 9- substituted tetracycline compound, comprising:

- 25 contacting a tetracycline compound with a nitrating agent, under conditions such that a nitro tetracycline compound is formed;
- contacting the nitro tetracycline compound with a hydrogenating agent, under conditions such that an amino tetracycline compound is formed; and
- contacting the amino tetracycline compound with an amino reactive
- 30 substrate, such that a 9- or 7- substituted tetracycline compound is formed.




- 80 -

87. The method of claim 86, wherein said substituted tetracycline compound is 9-substituted.
88. The method of claim 86, wherein said substituted tetracycline compound is 7-substituted.  
5
89. The method of claim 86, wherein the nitrating agent is  $\text{NaNO}_2$ .
90. The method of claim 86, wherein the nitrating agent is contacted with the tetracycline compound under acidic conditions.  
10
91. The method of claim 86, wherein said hydrogenating agent is hydrogen gas.
92. The method of claim 91, wherein said hydrogenating agent further comprises a transition metal catalyst.  
15
93. The method of claim 92, wherein said catalyst is platinum.
94. The method of claim 86, wherein said amino reactive compound is an isocyanate.  
20
95. The method of claim 86, wherein said amino reactive compound is isothiocyanate.
96. The method of claim 86, wherein said amino reactive compound is an unsubstituted or substituted chloroformate.  
25

- 81 -

97. A method for synthesizing a 7- or 9- substituted tetracycline compound of formula (I) comprising contacting a reactive intermediate with appropriate reagents under appropriate conditions, such that a substituted tetracycline compound is formed, wherein formula (I) is:



wherein:

X is  $\text{CHC}(\text{R}^{13}\text{Y}'\text{Y})$ ,  $\text{CR}^6\text{R}^6$ , S,  $\text{NR}^6$ , or O;

$\text{R}^2$  is hydrogen, alkyl, alkynyl, alkoxy, alkylthio, alkylsulfinyl, alkylsulfonyl, alkylamino, arylalkyl, aryl, heterocyclic, heteroaromatic or a prodrug moiety;

10

$\text{R}^4$  and  $\text{R}^{4'}$  are each hydrogen, alkyl, alkenyl, alkynyl, alkoxy, alkylthio, alkylsulfinyl, alkylsulfonyl, alkylamino, arylalkyl, aryl, heterocyclic, heteroaromatic or a prodrug moiety;

15  $\text{R}^{2'}$ ,  $\text{R}^3$ ,  $\text{R}^{10}$ ,  $\text{R}^{11}$  and  $\text{R}^{12}$  are each hydrogen or a pro-drug moiety;

$\text{R}^5$  is hydrogen, hydroxyl, or a prodrug moiety;

$\text{R}^6$ ,  $\text{R}^{6'}$ , and  $\text{R}^8$  are each independently hydrogen, alkyl, alkenyl, alkynyl, aryl, alkoxy, alkylthio, alkylsulfinyl, alkylsulfonyl, alkylamino, arylalkyl, or halogen;

$\text{R}^7$  is hydrogen, dialkylamino, heteroaryl-amino, or  $\text{NR}^{7c}\text{C}(=\text{W}')\text{WR}^{7a}$ ;

20  $\text{R}^{13}$  is hydrogen, hydroxy, alkyl; alkenyl; alkynyl; alkoxy; alkylthio; alkylsulfinyl; alkylsulfonyl; alkylamino; or an arylalkyl;

$\text{Y}'$  and  $\text{Y}$  are each independently hydrogen; halogen; hydroxyl; cyano, sulfhydryl; amino; alkyl; alkenyl; alkynyl; alkoxy; alkylthio; alkylsulfinyl; alkylsulfonyl; alkylamino; or an arylalkyl;

25  $\text{R}^9$  is hydrogen,  $\text{NR}^{9c}\text{C}(=\text{Z}')\text{ZR}^{9a}$ , or heteroaryl-amino;

Z is  $\text{CR}^{9d}\text{R}^{9e}$ ,  $\text{NR}^{9b}$ , or O;

$\text{Z}'$  is O or S;

$\text{R}^{9a}$ ,  $\text{R}^{9b}$ ,  $\text{R}^{9c}$ ,  $\text{R}^{9d}$ , and  $\text{R}^{9e}$  are each independently hydrogen, alkyl, alkenyl, alkynyl, alkoxy, alkylthio, alkylsulfinyl, alkylsulfonyl, arylsulfonyl,

- 82 -

alkoxycarbonyl, arylcarbonyl, alkylamino, arylalkyl, aryl, heterocyclic, heteroaromatic, absent, or a prodrug moiety;

W is  $\text{CR}^{7d}\text{R}^{7e}$ ,  $\text{NR}^{7b}$  or O;

W' is O or S; and

- 5  $\text{R}^{7a}$ ,  $\text{R}^{7b}$ ,  $\text{R}^{7c}$ ,  $\text{R}^{7d}$ , and  $\text{R}^{7e}$  are each independently hydrogen, alkyl, alkenyl, alkynyl, alkoxy, alkylthio, alkylsulfinyl, arylsulfonyl, alkoxycarbonyl, arylcarbonyl, alkylamino, arylalkyl, aryl, heterocyclic, heteroaromatic, absent, or a prodrug moiety;

- and pharmaceutically acceptable salts thereof, provided that  $\text{R}^9$  is not  
10 hydrogen when  $\text{R}^7$  is dialkylamino or hydrogen.


98. The method of claim 97, wherein said reactive intermediate is a 7- or 9-diazonium salt.

- 15 99. The method of claim 97, wherein said reactive intermediate is a 7- or 9-nitro compound.

100. The method of claim 97, wherein said reactive intermediate is a 7- or 9-thiourea.

- 20 101. The method of claim 97, wherein said reactive intermediate is a 7- or 9-thiocarboxamide.

102. A reactive intermediate, wherein said reactive intermediate is of the formula:



(I)

25

wherein:

X is  $\text{CHC}(\text{R}^{13}\text{Y}'\text{Y})$ ,  $\text{CHR}^6$ , S,  $\text{NR}^6$ , or O;

- 83 -

R<sup>2</sup> is hydrogen, alkyl, alkynyl, alkoxy, alkylthio, alkylsulfinyl, alkylsulfonyl, alkylamino, arylalkyl, aryl, heterocyclic, heteroaromatic or a prodrug moiety;

5 R<sup>4</sup> and R<sup>4'</sup> are each hydrogen, alkyl, alkenyl, alkynyl, alkoxy, alkylthio, alkylsulfinyl, alkylsulfonyl, alkylamino, arylalkyl, aryl, heterocyclic, heteroaromatic or a prodrug moiety;

R<sup>2'</sup>, R<sup>3</sup>, R<sup>10</sup>, R<sup>11</sup> and R<sup>12</sup> are each hydrogen or a pro-drug moiety;

R<sup>5</sup> is hydrogen, hydroxyl, or a prodrug moiety;

10 R<sup>6</sup> and R<sup>8</sup> are each independently hydrogen, alkyl, alkenyl, alkynyl, aryl, alkoxy, alkylthio, alkylsulfinyl, alkylsulfonyl, alkylamino, or an arylalkyl;

R<sup>13</sup> is hydrogen, hydroxy, alkyl; alkenyl; alkynyl; alkoxy; alkylthio; alkylsulfinyl; alkylsulfonyl; alkylamino; or an arylalkyl;

15 Y' and Y are each independently hydrogen; halogen; hydroxyl; cyano, sulfhydryl; amino; alkyl; alkenyl; alkynyl; alkoxy; alkylthio; alkylsulfinyl; alkylsulfonyl; alkylamino; or an arylalkyl;

R<sup>9</sup> is hydrogen, thiourea, diazonium salt, thiocarboxamide, or nitro;

R<sup>7</sup> is hydrogen, dialkylamino, thiourea, diazonium salt, thiocarboxamide, or nitro; and pharmaceutically acceptable salts thereof, provided that both R<sup>9</sup> is not hydrogen when R<sup>7</sup> is dialkylamino or hydrogen.

20

103. The reactive intermediate of claim 102, wherein R<sup>7</sup> is H, and R<sup>9</sup> is thiourea, diazonium salt, thiocarboxamide, or nitro moiety.

104. The reactive intermediate of claim 102, wherein R<sup>9</sup> is H, and R<sup>7</sup> is thiourea,  
25 diazonium salt, thiocarboxamide, or nitro moiety.

## INTERNATIONAL SEARCH REPORT

International Application No.

PCT/US 01/10342

## A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C07C271/30 C07C275/40 C07C271/58 C07C335/20 C07C235/84  
C07D277/42 A61K31/65

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C07C C07D A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, WPI Data, BEILSTEIN Data, CHEM ABS Data

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	<p>WO 96 34852 A (PFIZER ;VU CHI BAO (US)) 7 November 1996 (1996-11-07) page 1, line 18 -page 7, line 3 compounds of Tables 1 and 2, especially table 1, compounds 1-14</p> <p style="text-align: center;">--- -/--</p>	1-104



Further documents are listed in the continuation of box C.



Patent family members are listed in annex.

## \* Special categories of cited documents :

- \*A\* document defining the general state of the art which is not considered to be of particular relevance
- \*E\* earlier document but published on or after the international filing date
- \*L\* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- \*O\* document referring to an oral disclosure, use, exhibition or other means
- \*P\* document published prior to the international filing date but later than the priority date claimed

- \*T\* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- \*X\* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- \*Y\* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- \*Z\* document member of the same patent family

Date of the actual completion of the international search

29 August 2001

Date of mailing of the international search report

05/09/2001

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel (+31-70) 340-2040, Tx. 31 651 epo nl,  
Fax (+31-70) 340-3016

Authorized officer

O'Sullivan, P

## INTERNATIONAL SEARCH REPORT

International Application No.

PCT/US 01/10342

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	BARDEN T C ET AL: "GLYCYLCYCLINES 3. 9-AMINODOXYCYCLINECARBOXAMIDES" JOURNAL OF MEDICINAL CHEMISTRY, AMERICAN CHEMICAL SOCIETY. WASHINGTON, US, vol. 37, 30 September 1994 (1994-09-30), pages 3205-3211, XP000608350 ISSN: 0022-2623 page 3205, column 1, paragraph 2 scheme 1, compounds 4d,e,f page 3205, column 2, paragraph 2 scheme 2, compounds 9-18 page 3206, column 1, paragraph 1 scheme 3, compounds 34,35,39 ---	1-104
X	US 5 401 729 A (SUM PHAIK-ENG ET AL) 28 March 1995 (1995-03-28) column 1, line 26 - line 60 column 16, line 57 -column 17, line 30 examples, especially 3,4,6,26 ---	1-104
A	WO 99 37306 A (LEVY STUART B ;NELSON MARK L (US); TUFTS COLLEGE (US)) 29 July 1999 (1999-07-29) page 3, line 20 -page 4, line 14 examples 6,7 ---	1-104
A	US 4 024 272 A (ROGALSKI WERNER ET AL) 17 May 1977 (1977-05-17) cited in the application column 1, line 1 - line 36 claims -----	1-104

# INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US 01/10342

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9634852 A	07-11-1996	AU 693013 B	18-06-1998
		AU 5118696 A	21-11-1996
		CA 2219277 A	07-11-1996
		EP 0823893 A	18-02-1998
		JP 11504916 T	11-05-1999
		NZ 304095 A	28-10-1998
		US 6165999 A	26-12-2000
US 5401729 A	28-03-1995	US 5328902 A	12-07-1994
		AT 168992 T	15-08-1998
		AU 4461293 A	17-02-1994
		CA 2103836 A	14-02-1994
		CN 1083805 A,B	16-03-1994
		CZ 9301599 A	16-03-1994
		DE 69319976 D	03-09-1998
		DE 69319976 T	25-03-1999
		DK 582788 T	01-02-1999
		EP 0582788 A	16-02-1994
		ES 2119831 T	16-10-1998
		FI 933565 A	14-02-1994
		HK 1009598 A	20-04-2000
		HU 64943 A	28-03-1994
		HU 9500629 A	28-11-1995
		HU 74001 A,B	28-10-1996
		IL 106673 A	20-11-1997
		IL 119118 A	08-02-1998
		JP 6206855 A	26-07-1994
		LV 12358 A	20-10-1999
		LV 12358 B	20-01-2000
		MX 9304647 A	28-02-1994
		NO 932869 A	14-02-1994
		NZ 248357 A	26-04-1996
		PH 30007 A	29-10-1996
		PL 300063 A	21-02-1994
		PL 174115 B	30-06-1998
		PL 173904 B	29-05-1998
		RU 2125986 C	10-02-1999
		SG 47559 A	17-04-1998
		SK 85693 A	08-06-1994
		US 5380888 A	10-01-1995
		US 5466684 A	14-11-1995
		US 5495031 A	27-02-1996
		ZA 9305894 A	09-03-1994
WO 9937306 A	29-07-1999	AU 2334699 A	09-08-1999
		AU 2466099 A	09-08-1999
		EP 1047431 A	02-11-2000
		WO 9937307 A	29-07-1999
US 4024272 A	17-05-1977	DE 2442829 A	18-03-1976
		AT 344890 B	10-08-1978
		AT 689275 A	15-12-1977
		AU 8459975 A	10-03-1977
		BE 833048 A	03-03-1976
		CA 1068285 A	18-12-1979
		CS 191275 B	29-06-1979
		DD 121637 A	12-08-1976
		DK 397875 A	07-03-1976

# INTERNATIONAL SEARCH REPORT

Information on patent family members

Int: International Application No

PCT/US 01/10342

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
US 4024272 A		ES 440765 A	16-06-1977
		FR 2283690 A	02-04-1976
		GB 1467206 A	16-03-1977
		HU 173491 B	28-05-1979
		HU 176136 B	28-12-1980
		IE 41675 B	27-02-1980
		IL 48044 A	17-12-1978
		JP 51054564 A	13-05-1976
		LU 73332 A	13-04-1976
		NL 7510505 A	09-03-1976
		SE 7509880 A	08-03-1976
		ZA 7505699 A	25-08-1976

---